

Collaborative Staging Manual and Coding Instructions

Version 01.04.00



National
Cancer Institute
of Canada

Institut national
du cancer
du Canada

**COLLABORATIVE
STAGING MANUAL
AND
CODING INSTRUCTIONS**

Collaborative Staging Task Force
of the American Joint Committee on Cancer

Part I
Version 01.04.00

incorporating updates through October 31, 2007

NIH Publication Number 04-5496
U.S. Department of Health and Human Services
National Institutes of Health
National Cancer Institute

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

COPYRIGHT INFORMATION

All material in this manual is in the public domain and may be reproduced or copied without permission. However, citation as to source is requested.

SUGGESTED CITATION

Collaborative Staging Task Force of the American Joint Committee on Cancer. *Collaborative Staging Manual and Coding Instructions, version 01.04.00*. Jointly published by American Joint Committee on Cancer (Chicago, IL) and U.S. Department of Health and Human Services (Bethesda, MD), 2004. NIH Publication Number 04-5496. Incorporates updates through September 8, 2006.

Collaborative Staging (CS) Task Force Members (2006-2007)

<u>Member</u>	<u>Representing</u>
Stephen Edge, MD, Chair	American Joint Committee on Cancer
Cynthia Boudreaux, LPN, CTR	National Cancer Registrars Association
Gayle Clutter, RT, CTR	National Program of Cancer Registries, Centers for Disease Control and Prevention
Ingrid Friesen, HRT	Statistics Canada
April Fritz, BA, RHIT, CTR	Member at Large
Frederick L. Greene, MD, FACS	Ex-officio member, former Chair, AJCC
David L. Page, MD, FCAP	Chair, AJCC
Tom Rawson	National Program of Cancer Registries, Centers for Disease Control and Prevention
Lynn Ries, MS	Surveillance, Epidemiology, and End Results Program, National Cancer Institute
Colleen Sherman, RHIA, CTR	North American Association of Central Cancer Registries
Jennifer Seiffert, MLIS, CTR	National Program of Cancer Registries, Centers for Disease Control and Prevention
Valerie Vesich, RHIT, CTR	Member at Large

American Joint Committee on Cancer Staff

Connie Bura, Administrative Director, Cancer Programs, ACoS
Karen Pollitt, AJCC Administrator
Donna Gress, RHIT, CTR, AJCC Technical Specialist

Questions regarding this document and Collaborative Staging in general should be directed to:

Donna Gress, RHIT, CTR dgress@facs.org 312-202-5410
American Joint Committee on Cancer
633 North Saint Clair Street
Chicago, IL 60611

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

TABLE OF CONTENTS

PART I. INTRODUCTION AND GENERAL INSTRUCTIONS

INTRODUCTION	I-1
Changes in Abstracting Rules	I-2
How the Collaborative Staging System Works	I-4
Mapping and the Computer Algorithm	I-4
Table 1. Allowable Values and Format for Collaborative Staging Data Items	I-5
Figure 1. Schematic Diagram of Relationships of Inputs and Outputs for Collaborative Staging	I-7
How Mapping Was Determined	I-8
GENERAL INSTRUCTIONS	
General Instructions for Using the Collaborative Staging System:	I-11
Codes and Coding Instructions	
Table 2. Site Specific Factors Used For Primary Site/Histology Schemas	I-13
Structure and Format of Site/Histology-Specific Code Schemas	I-14
Coding “None” Vs. “Unknown” in the Collaborative Staging System, TNM and Summary Stage	I-14
Choosing the Correct Coding Schema for a Case	I-15
Table 3. Histology-specific Coding Schemas	I-15
Table 4. Schemas Where Tumor Size Is Required for AJCC Staging	I-16
Table 5. Schemas That Do Not Use Tumor Size for AJCC Staging	I-17
Table 6. Schemas for Which TNM Is Not Applicable	I-18
Death Certificate Only Cases	I-18
Use of Autopsy Information in Collaborative Staging	I-18
Definitions of Adjacent Tissues, Structures, and Organs	I-19
Ambiguous Terminology	I-20
How to Code the Collaborative Staging System Data Elements	I-21
INDIVIDUAL DATA FIELD CODING GUIDELINES	
CS Tumor Size	I-25
CS Extension	I-28
CS Tumor Size/Ext- Eval	I-30
CS Lymph Nodes	I-33
Coding Regional Lymph Nodes for Head and Neck Sites	I-35
Figure 2. Layout of Site-Specific Factors for Head and Neck Sites	I-36
Definitions of Levels for Head and Neck Sites	I-37
Coding Regional Nodes for Breast	I-38
CS Reg Nodes Eval	I-43
Regional Nodes Positive	I-45
Regional Nodes Examined	I-46
CS Mets at Dx	I-47
CS Mets Eval	I-49
CS Site-Specific Factor 1	I-51
CS Site-Specific Factor 2	I-53
CS Site-Specific Factor 3	I-55
CS Site-Specific Factor 4	I-57
CS Site-Specific Factor 5	I-59
CS Site-Specific Factor 6	I-61

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Table of Contents, continued

Additional Collaborative Staging Tables			I-63
Figure 3. Example of “Extension Size Table” for Breast Schema			I-64
APPENDICES			
Appendix 1.	Determining Descriptive Tumor Size		I-66
Appendix 2.	Output Values, Storage Codes, and Display String Description String		
	a. Allowable T Codes		I-67
	b. Allowable N Codes		I-68
	c. Allowable M Codes		I-69
	d. Allowable Stage Codes		I-69
	e. Allowable Summary Stage Codes		I-71
Appendix 3.	Summary Stage Conversion Algorithm for All Schemas		I-72
Appendix 4.	Site Specific Factors		I-76
Appendix 5.	Histology Exclusion Groups		I-81
INDEX TO PART I			I-84
 PART II. PRIMARY SITE SCHEMA			
HEAD AND NECK			
Lip, Upper	II-1	Colon	II-185
Lip, Lower	II-9	Rectosigmoid, Rectum	II-193
Lip, Other	II-17	Anus	II-199
Base of Tongue, Lingual Tonsil	II-25	Liver and Intrahepatic Bile Ducts	II-203
Anterior 2/3 of Tongue	II-33	Gallbladder	II-209
Gum, Upper	II-41	Extrahepatic Bile Ducts	II-213
Gum, Lower, and Retromolar Area	II-49	Ampulla of Vater	II-217
Gum, NOS	II-57	Other Biliary and Biliary, NOS	II-221
Floor of Mouth	II-65	Pancreas: Head	II-225
Hard Palate	II-73	Pancreas: Body and Tail	II-229
Soft Palate, Uvula	II-81	Pancreas: Other and Unspecified	II-233
Other Mouth	II-89	Other and III-Defined Digestive Organs	II-237
Cheek (Buccal) Mucosa, Vestibule	II-97		
Parotid Gland	II-105	RESPIRATORY AND INTRATHORACIC	
Submandibular Gland	II-113	Nasal Cavity	II-241
Other and Unspecified Major Salivary Glands	II-121	Middle Ear	II-249
Tonsil, Oropharynx	II-129	Maxillary Sinus	II-257
Anterior Surface of Epiglottis	II-137	Ethmoid Sinus	II-265
Nasopharynx	II-145	Accessory (Paranasal) Sinuses	II-273
Pyramiform Sinus, Hypopharynx	II-153	Larynx, Glottic	II-281
Pharynx, NOS, and Other III-Defined Oral Cavity	II-161	Larynx, Supraglottic	II-289
		Larynx, Subglottic	II-297
		Larynx, Overlapping Lesion or NOS	II-305
		Trachea	II-313
		Lung	II-317
		Heart, Mediastinum	II-325
		Pleura	II-329
		Other and III-Defined Respiratory Sites and Intrathoracic Organs	II-333
GASTROINTESTINAL			
Esophagus	II-167		
Stomach	II-173		
Small Intestine	II-179		

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Table of Contents, continued

MUSCULOSKELETAL		EYE	
Bone	II-337	Conjunctiva	II-479
Skin	II-341	Melanoma of Conjunctiva	II-483
Skin of Eyelid	II-345	Cornea, Retina, Choroid, Ciliary	
Melanoma of Skin	II-349	Body, Eyeball, Overlapping	
Mycosis Fungoides and Sezary		and Other Eye	II-487
Disease	II-357	Melanoma of Iris and Ciliary Body	II-491
Peripheral Nerves and Autonomic		Melanoma of Choroid	II-497
Nervous System; Connective,		Melanoma of Other Eye	II-501
Subcutaneous and Other		Lacrimal Gland	II-505
Soft Tissues	II-361	Orbit	II-509
Retroperitoneum, Peritoneum	II-367	Retinoblastoma	II-513
BREAST AND FEMALE GENITAL		CENTRAL NERVOUS	
Breast	II-371	Brain and Cerebral Meninges	II-519
Vulva	II-381	Other Parts of Central Nervous	
Vagina	II-385	System	II-523
Cervix	II-389		
Corpus	II-395	ENDOCRINE	
Ovary	II-401	Thyroid Gland	II-527
Fallopian Tube	II-407	Thymus, Adrenal Gland, and Other	
Broad and Round Ligaments,		Endocrine Glands	II-533
Parametrium and Other Adnexa	II-413		
Other and Unspecified Female		KAPOSI SARCOMA of All Sites	II-539
Genital Organs	II-417		
Placenta	II-421		
MALE GENITAL		HEMATOPOIETIC AND LYMPHOID	
Penis	II-427	Hodgkin and Non-Hodgkin	
Prostate	II-431	Lymphoma	II-543
Testis	II-443	Hematopoietic, Reticuloendothelial,	
Other and Unspecified Male		Immunoproliferative and	
Genital Organs	II-449	Myeloproliferative Neoplasms	II-549
Scrotum	II-453		
Kidney	II-457	OTHER	
Renal Pelvis, Ureter	II-461	Other and III-Defined Sites, Unknown	
Bladder	II-465	Primary Site	II-553
Urethra	II-471		
Paraurethral Gland, Overlapping		STANDARD TABLES	II-557
Lesion and Unspecified Urinary			
Organs	II-475		
INDEX OF SCHEMA BY ICD-O-3 PRIMARY SITE CODE			II-563
ALPHABETIC INDEX TO PART II: SITE SCHEMA			II-568

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Page left blank.

Collaborative Staging Manual and Coding Instructions Part I

General Instructions

INTRODUCTION

The Collaborative Staging Task Force, formed in 1998, was convened to address the issue of discrepancies in staging guidelines among the three major staging systems used in the United States. This project is sponsored by the American Joint Committee on Cancer (AJCC) in collaboration with the National Cancer Institute Surveillance, Epidemiology and End Results Program (NCI-SEER); Centers for Disease Control and Prevention National Program of Cancer Registries (CDC/NPCR); National Cancer Registrars Association (NCRA); North American Association of Central Cancer Registries (NAACCR); American College of Surgeons Commission on Cancer (CoC), and Canadian Cancer Society / National Cancer Institute of Canada (CCS-NCIC).

The initial focus of the Task Force was to develop a translation or other method of conversion between the TNM staging system of the AJCC and the SEER Summary Staging System. Such a translation would eliminate duplicate data collection by registrars reporting to clinical (facility-based) and epidemiologic (population-based central) registries, address the concerns of clinicians for more clinically relevant data as well as the public health sector's concerns about data reproducibility over time, and provide a higher degree of compatibility between the systems that would expand data-sharing opportunities.

The Collaborative Staging System is a carefully selected set of data items that describe how far a cancer has spread at the time of diagnosis. Most of the data items have traditionally been collected by cancer registries, including tumor size, extension, lymph node status, and metastatic status. New items were created to collect information necessary for the conversion algorithms, including the evaluation fields that describe how the collected data were determined, and site/histology-specific factors that are necessary to derive the final stage grouping for certain primary cancers. In addition to the items coded by the registrar, this unified data set also includes several data items derived from the computer algorithms that classify each case in multiple staging systems: the sixth edition of the AJCC TNM system (TNM), Summary Stage 1977 (SS77), and SEER Summary Stage 2000 (SS2000).

AJCC TNM staging provides forward flexibility and clinical utility for individual cancer cases. TNM is dynamic and is changed periodically to meet the decision-making needs of clinicians regarding appropriate treatment methods and the evaluation of their results. The AJCC TNM staging system uses three basic descriptors that are then grouped into stage categories. The first component is "T," which describes the extent of the primary tumor. The next component is "N," which describes the absence or presence and extent of regional lymph node metastasis. The third component is "M," which describes the absence or presence of distant metastasis. The final stage groupings (determined by the different permutations of "T," "N," and "M") range from Stage 0 through Stage IV. The stage group is generated when specific criteria are met in the TNM system, for example, prostate cancer stage grouping will only be generated for adenocarcinomas. When a case does not meet the criteria for stage grouping, the result will be reported as Not Applicable. An example of this type of case is leiomyosarcoma of the uterus, which is specifically excluded from TNM staging in both the uterus and the soft tissue sarcoma chapter. The Collaborative Staging System is based on, and compatible with, the terminology and staging in the sixth edition of the *AJCC Cancer Staging Manual*,¹ published in 2002. The general rules of the TNM system have been incorporated into the general rules for Collaborative Staging.

Summary Staging provides a measure for cancer surveillance with longitudinal stability for population-based cancer registries. Summary staging is a single digit system and has only eight categories: in situ, local, regional to lymph nodes, regional by direct extension, both regional lymph nodes and regional extension, regional not otherwise specified, distant, and unknown. It is less complex than other staging systems and was developed for registrars and epidemiologists who want some information on stage but did not wish to collect the more detailed EOD or TNM system. Summary Staging can be useful when a series of cases is so small that only general categories produce enough data for meaningful analysis. The

Collaborative Staging Manual and Coding Instructions Part I General Instructions

version of Summary Staging commonly used dates from 1977²; the site-specific sections were revised and updated in a new edition published in 2001³.

The Collaborative Staging System uses a modified EOD format to collect information about each case. The SEER Extent of Disease (EOD)⁴ coding system provided longitudinal stability for epidemiological and cancer control studies. More detailed than the Summary Staging System, EOD was developed to assure consistency over time as other staging systems changed. EOD also allows collected data to be collapsed into different and previous staging systems. SEER EOD is a five-field, 10 digit system: tumor size (3 digits), extension of the primary tumor (2 digits), regional lymph node involvement (highest specific lymph node chain involved by tumor) (1 digit), the number of pathologically reviewed regional lymph nodes that are positive (2 digits), and the number of pathologically examined regional lymph nodes (2 digits).

CHANGES IN ABSTRACTING RULES

Note: This introductory discussion refers to schemas based on primary site when in fact some schemas, such as melanoma and lymphoma, are based on histologic type. The schemas are referred to as site-specific for the sake of brevity.

Agreement among the participating organizations has resulted in resolution of the rule for timing of data collection and the development of standardized coding rules so that a single format can be used to collect stage information. The timing rule effective 1/1/2004 for Collaborative Staging is: "use all information gathered through completion of surgery(ies) in first course of treatment, or all information available within four months of the date of diagnosis in the absence of disease progression, whichever is *longer*." This timing rule change allows the CS Data Set to derive a "best stage" using pathologic data supplemented by clinical data.

Disease progression is defined as further direct extension or distant metastasis known to have developed after the diagnosis was established. Information about tumor extension, lymph node involvement, or distant metastasis obtained after disease progression is documented should be excluded from the Collaborative Staging fields. Collaborative Staging represents the aggregate information obtained during the period of diagnosis and work-up, not just the initial contact with the patient. For example, within the limits of the timing rule, if further diagnostic tests show more precise extension or a more precise tumor size, this revised information is not considered disease progression. In other words, Collaborative Staging does not consider as disease progression a change from lack of evidence of disease (status unknown) to known status of disease (negative or positive). However, a change from negative status to positive is disease progression. Take, for example, an asymptomatic patient who is treated surgically. She then develops bone pain and is found to have osseous metastases within a few weeks of surgery. This would be considered disease progression because she was asymptomatic at the time her treatment decisions were made. Furthermore, if the treatment plan is discontinued or changed due to a revised disease status, this is progression of disease and collection of Collaborative Staging information stops at this point.

Other rule modifications have been made and are printed in the site/histology-specific chapters.

In the process of bringing together the principles of Summary Stage, the TNM categories and stage groupings, and the SEER Extent of Disease coding structure, the Collaborative Staging System has also attempted to update abstracting rules to deal with the contemporary health care environment, in which completeness of staging documentation in the medical record has become an issue. In many circumstances, a patient's insurance will not pay for an imaging study or lab test that is expected to be negative but may otherwise be considered part of an 'ideal' cancer staging workup. Similarly, the content of clinician notes has changed over time to simply report any symptomatic, suspicious, or involved areas rather than chronicle every body part that is normal. This change in documentation is a source of

Collaborative Staging Manual and Coding Instructions Part I General Instructions

frustration to data collectors who rely on statements of normalcy or negativity to establish the boundaries of how far the cancer has spread.

When clinical practice changes and data collection guidelines do not, the completeness of the data is affected. The implementation of the Collaborative Staging System introduces a paradigm shift in the collection of information documenting the extent of disease, particularly in the collection of information about regional lymph nodes or distant metastases for primary sites not easily examined by palpation, observation, physical examination, or other clinical methods. These 'inaccessible' primary sites include (but are not limited to) bladder, kidney, prostate, esophagus, stomach, lung, liver, corpus uteri, and ovary.

The Collaborative Staging System allows data collectors to record regional lymph nodes as negative (based on clinical evaluation) rather than unknown when there is no mention of regional lymph node involvement in the physical examination, pre-treatment diagnostic testing or surgical exploration, and the patient receives what would be usual treatment to the primary site (treatment appropriate to the stage of disease as determined by the physician). The basis for this shift in the approach to information missing from the medical record is that typically the clinician reports positive findings and tends to remain silent on some or all negative findings. This new coding guideline also allows data collectors to record distant metastasis clinically as none rather than unknown (again, based on clinical evaluation) when the clinician proceeds with usual treatment of the primary site, since this action presumes the absence of distant metastasis that would otherwise change the treatment approach.

These guidelines apply primarily to localized or early (T1, T2) stage in the TNM system for inaccessible primary sites such as those mentioned previously. The code(s) for unknown information can and should be used in situations where there is reasonable doubt that the tumor is no longer localized. An example would be when there is clinical evidence that a prostate cancer has penetrated through the capsule into the surrounding tissues (regional direct extension/T3a) and regional lymph node involvement is not mentioned.

By coding regional lymph nodes as negative and/or coding distant metastasis as none rather than coding these fields as unknown, the Collaborative Staging System computer algorithms will be able to derive a stage group that includes the best information.

For accessible primary sites that can be observed, palpated or examined without instruments, such as breast, oral cavity, skin, salivary gland, thyroid, and other organs, there should be some description of the regional lymph nodes. A statement such as "remainder of examination negative" is sufficient to code regional lymph nodes as clinically negative.

In summary, the developers of the CS model believe that it will improve the quality of data being collected by the cancer registry community. Uniform rules and standardized training will make it easier for cancer registry personnel to complete staging tasks.

Collaborative Staging Manual and Coding Instructions Part I General Instructions

HOW THE COLLABORATIVE STAGING SYSTEM WORKS

For each cancer case, the data collector determines the site of origin or general histology for the cancer. The data items specific to that cancer site/histology are extracted from the medical record and coded in the Collaborative Staging System fields. When data collection is complete, the data collector activates the computer algorithms to derive the values for the items in the TNM system and Summary Stage (both 1977 and 2000). These algorithms are provided in portable platform-independent form by the Task Force. The classification or stage of each tumor is actually determined by the computer in a consistent and accurate manner (see Mapping and the Computer Algorithm, below).

Table 1 lists the individual Collaborative Staging data items, both input and derived, together with their NAACCR item number, length and other information, as published in the NAACCR Standards Volume II Version 10.1, Chapter X, Data Descriptor Table (revised November 2003).

Mapping and the Computer Algorithm

Once the data collector has coded all of the Collaborative Staging System elements for a case (the input values), the coded values are passed to a computer program that generates the correct stage for the case in three systems: AJCC TNM, 6th edition; SEER Summary Stage 1977; and SEER Summary Stage 2000. The program returns a set of values for the set of output items included in Table 1. A schematic diagram of the relationship between the inputs and outputs is shown in Figure 1.

The output values are returned as a set of numeric codes designed for storage in the computerized abstract. Each of the numeric codes is also provided with a display value, or English language character string showing the meaning of the code. For example, a returned value of 12 for T means T1a, and a 15 means T1b. Appendix 2 shows all of the output values and their display strings.

The computer algorithm that generates the stages is based on the values in the mapping columns for each of the Collaborative Staging System data elements. Mapping is provided from each code to the appropriate category in TNM and each summary stage. Some schemas require reference to two or more tables to determine the appropriate category. The mapping column either contains the category or a pointer to a further table where the category can be determined. Once each of the categories is determined, a further step is performed to generate the final stage groups. An example of the type of reference table used in this final step is shown in Appendix 3 for converting the results of the individual CS Extension, CS Lymph Nodes and CS Mets at Dx field to Summary Stage 1977 and Summary Stage 2000. For TNM stage grouping, the tables are schema-specific. Although the data collector does not code the stage groups directly, the rules by which the stages are derived are explicit in all of the tables, and the logic that the computer program follows should be fully evident from the tables available to the data collector.

As part of the output of the CS algorithm, two additional fields should be stored by the computer in the CS data base: CS Version 1st and CS Version Latest. CS Version 1st is the number of the version initially used to code CS fields and may be updated if cases are recoded, for example for a special study, using a later version of the Collaborative Staging manual. Depending on the structure of the registry software, CS Version 1st could be stored automatically by the computer or entered manually by the abstractor. The meaning and interpretation of CS Version 1st will be dependent on vendor implementation and local practices. This field should be interpreted with caution in a dataset where the actual coding procedures are unknown. CS Version Latest is the number of the version of the CS algorithm used most recently to derive the CS output fields and should be updated by the computer (rather than manually) every time the CS Derived items are re-computed.

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

OBSOLETE CODES

From time to time, it is necessary to revise Collaborative Staging (CS) coding tables by reassigning concepts from one code to another to maintain the underlying structure and rules for code assignment. This can occur when a single code needs to be split into more than one code, or when a structure needs to be move from one table to another (for example, a lymph node being moved from CS Lymph Nodes to CS Mets at Dx). Codes in CS tables will not be deleted while users have data coded with those codes. Instead, the codes will be marked as OBSOLETE in their descriptions, and instructions will be provided for handling previously coded data.

In some cases, it may be possible to perform global corrections on prior data without manual review. In other cases, such as when a code is being split, it will be necessary to manually review abstracts and recode them. Guidance for handling each instance of OBSOLETE will be provided when the change is published.

The designation of OBSOLETE is an official part of the description of the code, and it should be displayed to users, for example, in pick lists for coding new data so that the codes are not used into the future, and in translation of codes in displays or printouts of abstracts.

Table 1. Allowable Values and Format for Collaborative Staging Data Items						
INPUT ITEMS						
Data Item Name	NAACCR Data Item Number	Char-acter Length	Allowable Values (site-specific unless otherwise stated)	Right Justified, Zero filled	Blanks: Yes or No	NAACCR Ver 10.x Column #
CS Tumor Size	2800	3	000-999	Yes	No	629-631
CS Extension	2810	2	00-99	Yes	No	632-633
CS Tumor Size/Ext Eval	2820	1	0-9	N/A	No	634-634
CS Lymph Nodes	2830	2	00-99	Yes	No	635-636
CS Reg Nodes Eval	2840	1	0-9	N/A	No	637-637
Regional Nodes Examined	830	2	00-90, 95, 96, 97, 98, 99 (all sites)	Yes	No	541-542
Regional Nodes Positive	820	2	00-90, 95, 97, 98, 99 (all sites)	Yes	No	539-540
CS Mets At Dx	2850	2	00-99	Yes	No	638-639
CS Mets Eval	2860	1	0-9	N/A	No	640-640
CS Site-Specific Factor 1	2880	3	000-999	Yes	No	641-643
CS Site-Specific Factor 2	2890	3	000-999	Yes	No	644-646
CS Site-Specific Factor 3	2900	3	000-999	Yes	No	647-649
CS Site-Specific Factor 4	2910	3	000-999	Yes	No	650-652
CS Site-Specific Factor 5	2920	3	000-999	Yes	No	6535-655
CS Site-Specific Factor 6	2930	3	000-999	Yes	No	656-658


**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Table 1 continued. Allowable Values and Format for Collaborative Staging Data Items						
OUTPUT ITEMS						
Data Item Name	NAACCR Data Item Number	Character Length	Allowable Values	Right Justified, Zero filled	Blanks: Yes or No	NAACCR Ver 10.x Column #
Derived AJCC T	2940	2	00, 01, 05, 06, 07, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 29, 30, 31, 32, 33, 39, 40, 41, 42, 43, 44, 49, 88, 99	N/A	N/A	659-660
Derived AJCC N	2960	2	00, 01, 02, 03, 04, 09, 10, 11, 12, 13, 18, 19, 20, 21, 22, 23, 29, 30, 31, 32, 33, 39, 88, 99	N/A	N/A	662-663
Derived AJCC M	2980	2	00, 10, 11, 12, 13, 19, 88, 99	N/A	N/A	665-666
Derived AJCC T Descriptor	2950	1	c, p, a, y	N/A	N/A	661-661
Derived AJCC N Descriptor	2970	1	c, p, a, y	N/A	N/A	664-664
Derived AJCC M Descriptor	2990	1	c, p, a, y	N/A	N/A	667-667
Derived AJCC Stage Group	3000	2	00, 01, 02, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 70, 71, 72, 73, 74, 88, 90, 99	N/A	N/A	668-669
Derived AJCC Flag	3030	1	Blank, 1, 2	N/A	Yes	672-672
Derived SS1977	3010	1	Blank, 0, 1, 2, 3, 4, 5, 7, 8, 9	N/A	Yes	670-670
Derived SS1977 Flag	3040	1	Blank, 1, 2	N/A	Yes	673-673
Derived SS2000	3020	1	Blank, 0, 1, 2, 3, 4, 5, 7, 8, 9	N/A	Yes	671-671
Derived SS2000 Flag	3050	1	Blank, 1, 2	N/A	Yes	674-674
CS Version 1 st	2935	6	000000-999999	N/A	Yes	705-710
CS Version Latest	2936	6	000000-999999	N/A	Yes	711-716

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Figure 1

Schematic Diagram of Relationships of Inputs and Outputs for Collaborative Staging



Collaborative Staging Manual and Coding Instructions Part I General Instructions

How Mapping Was Determined

The Collaborative Staging Task Force based its codes for the extension, lymph nodes, and metastases fields on SEER's Extent of Disease, which had been designed to accommodate collapsing into TNM 3rd edition and the SEER Summary Stages. Some fundamental restructuring of the EOD codes was necessary to accommodate the sixth edition of TNM with its greater detail and supplementary prognostic information. For example, in EOD, all lymph node involvement (regional and distant) was coded in the lymph nodes field. In Collaborative Staging, regional lymph node involvement is coded in the CS lymph node field, and distant lymph node involvement is coded with other distant metastases. In each table, codes were added or combined where necessary to accommodate the 6th edition of TNM. The following rules and procedures were used to determine the correct mapping to TNM 6th edition:

- **Downstaging rule.** The Collaborative Staging Task Force applied the stated rule from the AJCC manual, "If there is doubt concerning the T, N, or M classification to which a particular case should be assigned, then the lower (less advanced) category should be assigned." When a mapping could be made to more than one classification, for example, T1 or T2, the mapping was always made to the lower or less extensive category. Occasionally this rule did not seem to apply, for example, when a lower category seemed to provide an exclusive list, while the higher category was more general. The downstaging rule was not applied to the assignment of stage group, only to the assignment of T, N, and M classification.
- **Use of NOS.** The Collaborative Staging Task Force added NOS (not otherwise specified) to some of its T, N, M, and stage group categories for clarity and ease of processing. The NOS is added when a further breakdown of the T, N, and M permutations into subsets is available, but the correct subset cannot be determined. NOS can appear in both the descriptions of codes and the mapping. This NOS terminology is not official AJCC usage. The NOS can safely be ignored in reports and analyses when it is not a useful distinction. In addition, the data collector should only code to a category such as "Stated as T1 NOS" when the appropriate subset (e.g., T1a or T1b) cannot be determined.

Example. For glottic larynx, T1 means "Tumor limited to the vocal cord(s) . . ." T1a means tumor limited to one vocal cord, and T1b means tumor involves both vocal cords. In Collaborative Staging, the subgroup of T1 NOS is designated for use when the tumor is known to be limited to the vocal cords, but it cannot be determined whether one or both cords are involved. In Collaborative Staging, the category T1 would be used to mean all of the T1's, including the T1a's, T1b's, and T1 NOS's.

REFERENCES

1. Greene FL, Page DL, Fleming ID et al. *AJCC Cancer Staging Manual, sixth edition*. American Joint Committee on Cancer. New York: Springer-Verlag, 2002.
2. Shambaugh EM and Weiss MA. *SEER Summary Staging Guide 1977*. Bethesda, MD: National Cancer Institute, NIH Publication Number 98-2313, reprinted December 1997.
3. Young JL, Roffers SD, Ries LAG, Fritz AG, Hurlbut AA (eds). *Summary Staging Manual 2000: Codes and Coding Instructions*. Bethesda, MD: National Cancer Institute, NIH Publication Number 01-4969, 2001
4. Fritz AG and Ries LAG (eds). *SEER Extent of Disease Coding, 1998: Codes and Coding Instructions, Third Edition*. Bethesda, MD: National Cancer Institute, NIH Publication Number 98-1999, April 1998.

General Rules and Instructions

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Page left blank.

**General Instructions
for Using the Collaborative Staging System
Codes and Coding Instructions**

The Collaborative Staging System schemas consist of the 15 data fields necessary to derive T, N, M, and Stage Group according to the sixth edition of the *AJCC Cancer Staging Manual*; Summary Stage 1977; and SEER Summary Stage 2000.

This manual provides codes and coding instructions for the process of data entry. In order to derive the desired T, N, M, and Stage Group in the TNM system or the Summary Stage(s), the computer algorithms described in the introduction must be used. This manual provides the logic of the computer algorithms in table format for each schema, but is not intended to be used for generating the stages manually, because for some sites, additional tables are necessary to determine T, N, M, or Stage Group. These additional tables are available for review on the Collaborative Staging web site, <http://www.cancerstaging.org>

These schemas apply to cases diagnosed January 1, 2004 and later. **Do NOT use these schemas for cases diagnosed prior to January 1, 2004**; cases diagnosed prior to 01/01/2004 should be coded to whatever coding system was in effect at the time of diagnosis.

General Guidelines

Note: These general instructions refer to schemas based on primary site when, in fact, some schemas, such as melanoma and lymphoma, are based on histologic type. The schemas are referred to as site-specific for the sake of brevity.

1. Collaborative Staging is collected on all cases regardless of whether they are microscopically confirmed. A description of the type of diagnostic confirmation is collected in a separate data item. The diagnostic confirmation field can be used to exclude non-microscopically confirmed cases during analysis as necessary, since the *AJCC Cancer Staging Manual* states that “all cases should be microscopically confirmed. Cases not microscopically confirmed should be coded from the schema for the site/histology the clinician considers most likely to be the primary.”
2. Collaborative Staging is collected on all sites/histologies. Summary Stage 1977 and Summary Stage 2000 are generated for all sites and histologies. The TNM elements and stage group are only generated for cases that meet the TNM criteria. For example, there is no TNM schema for brain.
 - a. The Collaborative Staging System consists of 94 schemas, most of which are site-specific. Some malignancies that can develop in many parts of the body are coded according to the histology of the case. For example, all lymphomas are coded according to the lymphoma schema, regardless of the organ in which the lymphoma develops.
3. All schemas apply to all histologies unless otherwise noted. Summary Stage 1977 and Summary Stage 2000 are generated for all histologies. The computer algorithms for determining the final TNM stage group take into account any histologies that are excluded from TNM staging. For example, the TNM schema for prostate applies only to adenocarcinomas. For excluded histologies, the computer algorithm returns values representing “Not Applicable,” meaning that AJCC T, N, M, and Stage Group are not generated for that site-histology combination.

Collaborative Staging Manual and Coding Instructions Part I
General Instructions

4. **Timing of Data Collection:** The data collected in the Collaborative Staging System are limited to
 - information gathered through completion of surgery(ies) in first course of treatment, OR
 - all information available within four months of the date of diagnosis in the absence of disease progression (metastasis known to have developed after the diagnosis was established should be excluded)
 - whichever is *longer*.
5. Site-specific and histology-specific guidelines take precedence over general guidelines. Always read the notes pertaining to a specific site or histology schema.
6. For each field, code the highest applicable number. (Exception: codes for Unknown, Not Applicable, and NOS categories such as Localized, NOS do not take priority over more specific codes with lower numbers.) The codes are ordered in a hierarchy so that increasing numbers generally indicate increasing degrees of tumor involvement. The hierarchies are not the same for the different staging systems, and Collaborative Staging generally follows the hierarchies of the TNM system.
 - a. Combination codes (for example, code 35 for “25 plus 30”) have been assigned when using the higher number does not result in the appropriate mapping for all three stage groups. Combination codes have been omitted when use of a higher number results in correct mapping for all three staging systems.
7. For the fields CS Tumor Size, CS Extension, CS Lymph Nodes, and CS Mets at DX, Collaborative Staging records the greatest extent of disease based on combined clinical and operative/pathological assessment.
 - a. Gross observations at surgery are particularly important when all malignant tissue is not removed. In the event of a discrepancy between pathology and operative reports concerning excised tissue, priority is given to the pathology report.
 - b. Clinical information, such as a description of skin involvement for breast cancer and size of the primary lesion and distant lymph nodes for any site, can change the stage. Clinical information should be reviewed carefully to assure accurate recording of the Collaborative Staging data set.
8. When the patient does not receive preoperative treatment and the operative/pathology information disproves the clinical information, code the operative/pathology information.
9. When the patient does receive preoperative treatment, the greatest extent of disease prior to the beginning of treatment should be recorded. Preoperative, or neoadjuvant, treatment is defined as systemic (chemotherapy, hormone therapy, or immunotherapy) treatment or radiation therapy that is administered as an attempt to shrink the tumor, improve resectability, or control symptoms before the patient undergoes surgery. In the infrequent situation where post-operative disease is more extensive despite neoadjuvant treatment, this can be coded in the method of evaluation field for extension, regional lymph nodes or metastases at diagnosis.
10. The fields Reg LN Pos and Reg LN Exam are based on pathologic (microscopic) information only.
11. The fields CS Tumor Size/Ext Eval, CS Reg Nodes Eval, and CS Mets Eval document how the most extensive tumor was established as well as whether the patient received preoperative treatment.
12. Site-Specific Factors (SSFs) are included in every schema. They are incorporated into the staging algorithms when additional information is necessary to derive tumor (T), lymph node (N), metastasis (M), or TNM stage group, or where the factor is considered to be of clinical or prognostic importance. Information formerly coded as tumor markers, such as estrogen receptor assay or progesterone receptor assay for breast, is coded in site-specific factors. For sites/histologies where some or all site specific factors are not used, they are coded 888, not applicable. Table 2 lists the schemas that require one or more Site Specific Factors. Appendix 4 lists the names of each site specific factor by schema.

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Table 2. Site Specific Factors Used For Primary Site/Histology Schemas

SSF	Sites/histologies where used		
1	head and neck* colon rectum liver pleura melanoma	mycosis fungoides breast ovary placenta prostate testis melanoma/conjunctiva melanoma/choroid	melanoma/iris and ciliary body retinoblastoma brain other cns thyroid other endocrine Kaposi sarcoma lymphoma
2	head and neck*, liver, melanoma, breast, prostate, testis, lymphoma		
3	head and neck*, melanoma, breast, prostate, testis, lymphoma		
4	head and neck*, melanoma, breast, prostate, testis		
5	head and neck*, breast, prostate, testis		
6	head and neck*, breast, prostate		

* head and neck includes the following schemas: upper lip; lower lip; other lip; base of tongue; anterior tongue; upper gum; lower gum and retromolar trigone; other gum; floor of mouth; hard palate; soft palate/uvula; other mouth; buccal mucosa; parotid gland; submandibular gland; other salivary glands; oropharynx; anterior surface of epiglottis; nasopharynx; pyriform sinus/hypopharynx; other pharynx; nasal cavity; middle ear; maxillary sinus; ethmoid sinus; other sinus; glottic larynx; supraglottic larynx; subglottic larynx; other larynx

-
13. Metastasis known to have developed after the initial extent of disease was established (in other words, disease progression) should be excluded when determining the farthest extent of disease at the time of diagnosis.
 14. Autopsy reports are used in coding the Collaborative Staging System in the same way as are pathology reports, applying the same rules for inclusion and exclusion.
 15. The extent of disease may be described only in terms of T (tumor), N (node), and M (metastasis) characteristics. In such cases, assign the code in the appropriate field that corresponds to the TNM information. If there is a discrepancy between documentation in the medical record and the physician's assignment of TNM, the documentation takes precedence. Cases of this type should be discussed with the physician who assigned the TNM.

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

STRUCTURE AND FORMAT OF SITE/HISTOLOGY-SPECIFIC CODE SCHEMAS

The schemas in this manual are listed according to the order of the first ICD-O-3 primary site code to which a schema applies. Schemas for which there is no TNM classification are included in ICD-O-3 sequence in the manual. Some of the histology-based schemas appear in site code order (for example, melanoma of the skin is with other skin schemas), and others are at the end of the list. Two indices to the schemas are provided at the end of this manual, one by ICD-O-3 code and the other by common primary site and histology terms.

Within the schemas themselves, the code structures for the various organs, lymph nodes, and other tissues are organized according to the T, N, and M categories (T1, then T2, then T3, for example). As such, they may not be sequential for Summary Stage definitions. Regardless of the relative order of the codes in the schemas, the staging algorithms will properly account for the information.

The categories of TNM are the basis for the CS Extension, CS Lymph Nodes and CS Mets at DX fields. Tissues categorized under T in the TNM system are listed in CS Extension and tissues categorized under M are listed in the CS Mets at DX field. However, for the Summary Staging (1977 and/or 2000) algorithms, there may be codes in the CS Extension field that map to regional direct extension or distant stage, and there may be codes in CS Mets at DX that map to regional or even localized disease. The details of the case should be coded in the fields where they are listed; the computer algorithm is designed to generate the correct stage. It should also be noted that information in fields other than CS Extension may be used to derive the T, N, M and Stage Group, for example tumor size and various site-specific factors.

CODING “NONE” VS. “UNKNOWN” IN THE COLLABORATIVE STAGING SYSTEM, TNM AND SUMMARY STAGE

As noted in the introduction, cancers of certain primary sites are not easily examined by palpation, observation, physical examination, or other clinical methods. These ‘inaccessible’ primary sites include, but are not limited to, bladder, kidney, prostate, esophagus, stomach, lung, liver, corpus uteri and ovary.

A new coding rule in the Collaborative Staging System applies to these inaccessible sites, primarily for localized or early (T1, T2) stage cancers. The Collaborative Staging System allows data collectors to record regional lymph nodes as negative (based on clinical evaluation) rather than unknown when there is no mention of regional lymph node involvement in the physical examination, pre-treatment diagnostic testing or surgical exploration, and the patient receives what would be usual treatment to the primary site (treatment appropriate to the stage of disease as determined by the physician).

This new coding guideline also permits data collectors to record distant metastasis clinically as none rather than unknown (again, based on clinical evaluation) when the clinician proceeds with usual treatment of the primary site, since this action presumes that there are no distant metastasis that would otherwise change the treatment approach.

The code(s) for unknown information can and should be used in situations where there is reasonable doubt that the tumor is no longer localized. For example, when there is clinical evidence that a prostate cancer has penetrated through the capsule into the surrounding tissues (regional direct extension/T3a) and regional lymph node involvement is not mentioned, it would be correct to code lymph node involvement and metastases at diagnosis as unknown in the absence of any specific information regarding nodes or distant metastases.

For accessible primary sites that can be observed, palpated or examined without instruments, such as breast, oral cavity, skin, salivary gland, thyroid, and other organs, there should be some description of the

Collaborative Staging Manual and Coding Instructions Part I General Instructions

regional lymph node status. A statement such as “remainder of examination negative” is sufficient to code regional lymph nodes as clinically negative.

CHOOSING THE CORRECT CODING SCHEMA FOR A CASE

Most of the Collaborative Staging System schemas apply to cases defined by their primary site codes in ICD-O-3. A few of the schemas apply to cases defined by their histologic type codes in ICD-O-3, and these schemas take precedence over the schema for the site. The histologically defined schemas are shown in Table 3.

TABLE 3. HISTOLOGY-SPECIFIC CODING SCHEMAS

Melanoma (ICD-O-3 morphology codes 8720-8790)
Kaposi sarcoma (9140)
Retinoblastoma (9510-9514)
Lymphoma (9590-9699 and 9702-9729)
Mycosis Fungoides (9700-9701)
Hematopoietic and reticuloendothelial system (9731-9989)

A case with one of these ICD-O-3 histologic types must be coded using the schema for the histologic type group.

Melanomas are further broken down by primary site code, as follows:

Malignant melanoma of the skin, vulva, penis and scrotum (C44.0-C44.9, C51.0-C51.2, C51.8-C51.9, C60.0-C60.1, C60.8-C60.9, C63.2)
Malignant melanoma of conjunctiva (C69.0)
Malignant melanoma of iris and ciliary body (C69.4)
Malignant melanoma of choroid (C69.3)
Malignant melanoma of other eye (C69.1, C69.2, C69.5, C69.8-C69.9)

For cases with all other histologic types, the correct schema to use is determined by the primary site code.

Each schema clearly states the applicable primary site codes and histologic type codes at the beginning of the schema.

Note: The appropriate site or histology schema to use for coding surgical treatment(s) may be different from the site or histology schema used for coding the Collaborative Staging data set. For example, an extralymphatic lymphoma of the stomach treated surgically would use the lymphoma schema in this manual to code Collaborative Staging, but surgery would be coded using the stomach codes for surgery of primary site. Refer to the treatment coding rules in the SEER Program coding manual or the FORDS manual for more details.

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Schemas Where Tumor Size is Necessary for AJCC Staging

In order to classify the T category for certain sites/histologies, it is necessary to know the size of the primary tumor, usually for T1 - T3. For the following sites/histologies, the size of the primary tumor must be recorded in order to assign the T category and derive a stage group. Tumor size is not necessary to assign Summary Stage. The name of the Collaborative Staging schema and its website file name (shown in parentheses) are double indented under the **TNM chapter** and *subsite* names. (See Table 4.)

TABLE 4. SCHEMAS WHERE TUMOR SIZE IS NECESSARY FOR AJCC STAGING

Lip and oral cavity

Lip

- Upper Lip (LipUpper)
- Lower Lip (LipLower)
- Other Lip (OthLip)

Oral Cavity

- Anterior Tongue (AntTongue)
- Upper Gum (GumUpper)
- Lower Gum (GumLower)
- Other Gum (OthGum)
- Floor of Mouth (FOM)
- Hard Palate (HardPalate)
- Buccal Mucosa (BuccalMucosa)
- Other Mouth (OthMouth)

Pharynx

Oropharynx

- Oropharynx (Oropharynx)
- Base of Tongue (BaseTongue)
- Soft Palate (SoftPalate)

Hypopharynx

- Hypopharynx (Hypopharynx)

Major Salivary Glands

- Parotid Gland (ParotidGland)
- Submandibular Gland
(SubmandibularGland)
- Other Salivary Gland (OthSalivary)

Thyroid

- Thyroid (Thyroid)

Anal Canal

- Anus (Anus)

Liver including Intrahepatic Bile Ducts

- Liver and intrahepatic bile ducts (Liver)

Exocrine Pancreas

- Pancreas Head (PancreasHead)
- Pancreas Body and Tail
(PancreasBodyTail)
- Other Pancreas (OthPancreas)

Lung

- Lung (Lung)

Bone

- Bone (Bone)

Soft tissue sarcoma

- Heart and Mediastinum
(HeartMediastinum)
- Soft Tissue (SoftTissue)
- Peritoneum (Peritoneum)

Carcinoma of the Skin

- Skin, Vulva, Penis, Scrotum–Carcinoma
(Skin)

Carcinoma of the Eyelid

- Skin of Eyelid–Carcinoma (SkinEyelid)

Breast

- Breast (Breast)

Vulva

- Vulva (Vulva)

Cervix Uteri

- Cervix (Cervix)

Kidney

- Kidney (Kidney)

Carcinoma of the Conjunctiva

- Conjunctiva–Carcinoma (Conjunctiva)

Malignant Melanoma of the Uvea

- Iris and Ciliary Body–Melanoma (*ciliary body only*) (MelanomaIrisCiliary)
- Choroid–Melanoma (MelanomaChoroid)

Carcinoma of the Lacrimal Gland

- Lacrimal gland–Carcinoma
(LacrimalGland)

Sarcoma of the Orbit

- Orbit (Orbit)

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Schemas That Do Not Use Tumor Size for AJCC Staging

In order to classify both summary stage and the AJCC T category for certain sites/histologies, it is necessary to know how far the tumor has extended in a contiguous, continuous or direct manner from its point of origin. For the following sites/histologies, the extension of the primary tumor must be recorded in order to assign the T category and derive a stage group. The name of the Collaborative Staging schema and its website file name (in parentheses) are double indented under the **TNM chapter** and *subsite* names. (See Table 5.)

TABLE 5. SCHEMAS THAT DO NOT USE TUMOR SIZE FOR AJCC STAGING

Pharynx	Melanoma of the Skin
<i>Nasopharynx</i>	Skin, Vulva, Penis, Scrotum–Melanoma
Nasopharynx (Nasopharynx)	(Melanoma)
Larynx	Vagina
Other Larynx (OthLarynx)	Vagina (Vagina)
<i>Glottic Larynx</i>	Corpus uteri
Glottic Larynx (GlotticLarynx)	Corpus uteri (Corpus)
<i>Supraglottic Larynx</i>	Ovary
Supraglottic Larynx (SupraLarynx)	Ovary (Ovary)
Anterior Surface of Epiglottis	Fallopian Tube
(AntEpiglottis)	Fallopian tube (FallopianTube)
<i>Subglottic Larynx</i>	Gestational trophoblastic tumor
Subglottic Larynx (SubLarynx)	Placenta (Placenta)
Nasal Cavity and Paranasal Sinuses	Penis
Nasal Cavity (NasalCavity)	Penis (Penis)
Maxillary Sinus (MaxillarySinus)	Prostate
Ethmoid Sinus (EthmoidSinus)	Prostate (Prostate)
Esophagus	Testis
Esophagus (Esophagus)	Testis (Testis)
Stomach	Renal Pelvis and Ureter
Stomach (Stomach)	Renal Pelvis and Ureter (RenalPelvis)
Small Intestine	Urinary Bladder
Small intestine (SmallIntestine)	Bladder (Bladder)
Colon and rectum	Urethra
Colon (Colon)	Urethra (Urethra)
Rectum (Rectum)	Malignant Melanoma of the Conjunctiva
Gallbladder	Conjunctiva–Melanoma
Gallbladder (Gallbladder)	(MelanomaConjunctiva)
Extrahepatic bile ducts	Malignant Melanoma of the Uvea
Extrahepatic bile ducts	Iris and Ciliary Body–Melanoma (<i>iris</i>
(ExtraHepaticDucts)	<i>only</i>) (MelanomaIrisCiliary)
Other Biliary and Biliary, NOS	Retinoblastoma
(OthBiliary)	Retinoblastoma (Retinoblastoma)
Ampulla of Vater	Lymphoid neoplasms
Ampulla (Ampulla)	Mycosis Fungoides (MF)
Pleural mesothelioma	Malignant Lymphoma (Lymphoma)
Pleura (Pleura)	

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

TABLE 6. SCHEMAS FOR WHICH AJCC STAGING IS NOT APPLICABLE

For the following schemas, TNM is not applicable. The name of the Collaborative Staging schema and its website file name (in parentheses) are shown below.

Other pharynx (OthPharynx)	Other endocrine (OthEndocrine)
Other digestive (OthDigestive)	Other eye (OthEye)
Middle ear (MiddleEar)	Melanoma of Other Eye (MelanomaOthEye)
Other sinus (OthSinus)	Kaposi sarcoma (KS)
Trachea (Trachea)	Hematopoietic, Reticuloendothelial,
Other respiratory (OthRespiratory)	Immunoproliferative and
Other adnexa (OthAdnexa)	Myeloproliferative Neoplasms
Other female genital (OthFemaleGen)	(HemeRetic)
Other male genital (OthMaleGen)	Other Ill-defined and Unknown Primary
Other urinary (OthUrinary)	Sites (OthIllDef)
Brain (Brain)	
Other CNS (OthCNS)	

DEATH CERTIFICATE ONLY CASES

Death Certificate **only** cases are coded as unknown (usually 9, 99, 999, etc.) or not applicable (usually 8, 88, 888, etc.) in all Collaborative Staging fields. Although there may be some site/histology-specific exceptions, the usual pattern for coding Death Certificate Only cases is as follows:

CS Tumor Size	999	CS Site-Specific Factor 1	888
CS Extension	99	CS Site-Specific Factor 2	888
CS Tumor Size/Ext Eval	9	CS Site-Specific Factor 3	888
CS Lymph Nodes	99	CS Site-Specific Factor 4	888
CS Reg Nodes Eval	9	CS Site-Specific Factor 5	888
Reg LN Pos	99	CS Site-Specific Factor 6	888
Reg LN Exam	99		
CS Mets at DX	99		
CS Mets Eval	9		

USE OF AUTOPSY INFORMATION IN COLLABORATIVE STAGING

Information obtained from autopsy may be used in either of two ways in the Collaborative Staging System. The evaluation fields must then be coded correctly to indicate how the autopsy information is to be interpreted. If a patient with a suspected diagnosis of cancer dies and an autopsy is performed, extent of disease information obtained from the autopsy may be included along with other clinical and pathologic information, if it meets the timing rules for inclusion. In this case, the computer algorithm will assign the T, N, or M to “p” (pathologic) classification. If cancer is not suspected at the time of autopsy, the extent of disease information from the autopsy is included, but the algorithm will assign the T, N, and M to the autopsy (a) classification of the TNM system rather than to clinical or pathologic evaluation. Each of the evaluation field schemas has appropriate codes to allow this distinction.

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

DEFINITIONS OF ADJACENT TISSUES, STRUCTURES, AND ORGANS

Adjacent connective tissue

Some of the Collaborative Staging System schemas for ill-defined or non-specific sites in this manual contain a code for adjacent connective tissue, which is defined here as the unnamed tissues that immediately surround an organ or structure containing a primary cancer. Use this code when a tumor has invaded past the outer border (capsule, serosa, or other edge) of the primary organ into the organ's surrounding supportive structures but has not invaded into larger structures or adjacent organs.

The structures identified in ICD-O-3 as connective tissue include the following: adipose tissue; aponeuroses; arteries; blood vessels; bursa; connective tissue, NOS; fascia; fatty tissue; fibrous tissue; ganglia; ligaments; lymphatic channels (not nodes); muscle; nerves (spinal, sympathetic and peripheral); skeletal muscle; subcutaneous tissue; synovia; tendons; tendon sheaths; veins; and vessels, NOS. In general, these tissues do not have specific names. These tissues form the framework of many organs, provide support to hold organs in place, bind tissues and organs together, and serve as storage sites for nutrients. Blood, cartilage and bone are sometimes considered connective tissues, but in this manual they would be listed separately.

Adjacent organs

Organs are anatomic structures with specific physiologic functions other than (or in addition to) support and storage. Continuous tumor growth from one organ into an organ anatomically next to the primary would be coded to the appropriate code for 'adjacent organs/structures' in the Collaborative Staging schemas for ill-defined and non-specific sites.

Adjacent structures

Connective tissues large enough to be given a specific name would be considered adjacent structures. For example, the brachial artery has a name, as does the broad ligament. Continuous tumor growth from one organ into an adjacent named structure would be coded to the appropriate code for 'adjacent organs/structures' in the Collaborative Staging for ill-defined or non-specific sites.

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

AMBIGUOUS TERMINOLOGY

Interpreting Ambiguous Terminology for Collaborative Staging

Determination of the cancer stage is both a subjective and objective assessment of how far the cancer has spread. Sometimes the clinician is hesitant to commit to a definite statement that a particular organ or tissue is involved by the cancer and uses what data collectors refer to as “ambiguous terminology.” The following lists can generally be used to interpret the intent of the clinician; however, if individual clinicians use these terms differently, the clinician’s definitions and choice of therapy should be recognized. If a term used in a diagnostic statement is not listed below, consult the clinician to determine the intent of the statement.

Consider as involvement

adherent
apparent(ly)
appears to
comparable with
compatible with
consistent with
contiguous/continuous with
encroaching upon*
extension to, into, onto, out onto
features of
fixation to another structure**
fixed**
impending perforation of
impinging upon
impose/imposing on
incipient invasion
induration
infringe/infringing
into*
intrude
invasion to into, onto, out onto
most likely
onto*
overstep
presumed
probable
protruding into (unless encapsulated)
suspected
suspicious
to*
up to

DO NOT Consider as Involvement

abuts
approaching
approximates
attached
cannot be excluded/ruled out
efface/effacing/effacement
encased/encasing
encompass(ed)
entrapped
equivocal
extension to without invasion/
involvement of
kiss/kissing
matted (except for lymph nodes)
possible
questionable
reaching
rule out
suggests
very close to
worrisome

* interpreted as involvement whether the description is clinical or operative/
pathological

** interpreted as involvement of other organ or tissue

Collaborative Staging Manual and Coding Instructions Part I
General Instructions

HOW TO CODE THE COLLABORATIVE STAGING SYSTEM DATA ELEMENTS

A one page summary of how to code using this manual

Note: This procedure focuses on only the Collaborative Staging data fields and assumes other registry operations such as case finding, completion of text fields and other data fields, edit checking and case submission are also being performed appropriately.

1. Before you begin to code using the Collaborative Staging System, read completely the general rules in this manual.
2. Read the medical record carefully to determine the primary site and histology and identify the correct ICD-O-3 codes. While you are reviewing the record, make mental notes about the tissues and lymph nodes that are involved by tumor.
3. If the histology is melanoma (8720-8790), Kaposi sarcoma (9140), retinoblastoma (9510-9514), lymphoma (9590-9699 and 9702-9729), mycosis fungoides (9700-9701), or hematopoietic and reticuloendothelial system (9731-9989), use the histology-specific schema for the appropriate histology-site combination.
4. Otherwise, turn to the correct site-specific schema in the Part II of this manual. Schemas are in ICD-O-3 order by the first code that uses the schema. Verify that you are in the correct chapter by confirming that the code is in the list at the beginning of the schema.
5. Begin assigning codes for the 15 fields in the Collaborative Staging System. Be sure to read the notes and follow the site/histology-specific instructions at the beginning of each data field. Some schemas may have site-specific factors associated with extension, lymph nodes or metastasis; keep these in mind as you assign the codes.
 - a. Code the tumor size in the CS Tumor Size field.
 - b. Code how far the tumor has directly spread in the CS Extension field.
 - c. Code how the farthest tumor spread was determined in the CS Tumor Size/Ext Eval field.
 - d. Code whether regional lymph nodes are involved in the CS Lymph Nodes field.
 - e. Code how the farthest regional node spread was determined in the CS Reg Node Eval field.
 - f. Code the number of positive regional lymph nodes from the pathology report in the Reg Nodes Pos field.
 - g. Code the number of regional lymph nodes examined by the pathologist in the Reg Nodes Exam field.
 - h. Code the farthest distant metastasis (including distant lymph nodes) in the CS Mets at Dx field.
 - i. Code how the distant metastasis was determined in the CS Mets Eval field.
 - j. Code the six site-specific factors. If the first site-specific factor is listed as "Not Applicable," code 888 in all site specific factors. Otherwise, code the specific information requested for each site specific factor. When the next site-specific factor is 888 Not Applicable, all the remaining site-specific factors will also be 888.

Congratulations! You have collected all the facts about the case and the codes are ready for the computer to convert into the T, N, M, Stage Group, Summary Stage 1977 and Summary Stage 2000. Depending on your software system, the final stage information may be derived now, when the case is saved, or prior to exiting the case. Finish the rest of the abstract, edit check it and save it.

When the computer derives the final stage information, the program will check the histology code and other coded information to determine whether T, N, M and Stage Group will be generated for the case. If the histology code is on the computer's exceptions list for that site, the T, N, M, and Stage Group will be reported as "Not Applicable." Summary Stage is generated for every case. The computer algorithm will also record which version of the Collaborative Staging System was used to derive the final stages.

**Collaborative Staging Manual and Coding Instructions Part I
General Instructions**

Page left blank.

**Coding Instructions for
Collaborative Staging Data Elements**

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

Page left blank.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS TUMOR SIZE

Item Length: 3

NAACCR Item #2800

Description

Records the largest dimension or diameter of the **primary tumor**, and is always recorded in millimeters. To convert centimeters to millimeters, multiply the dimension by 10. If tumor size is given in tenths of millimeters, record size as 001 if largest dimension or diameter of tumor is between 0.1 and 0.9 mm.

Code	Description
000	Indicates no mass or no tumor found; for example, when a tumor of a stated primary site is not found, but the tumor has metastasized.
001-988	Exact size in millimeters.
989	989 millimeters or larger.
990	Microscopic focus or foci only; no size of focus is given.
991	Described as "less than 1 cm"
992	Described as "less than 2 cm," or "greater than 1 cm," or "between 1 cm and 2 cm"
993	Described as "less than 3 cm," or "greater than 2 cm," or "between 2 cm and 3 cm"
994	Described as "less than 4 cm," or "greater than 3 cm," or "between 3 cm and 4 cm"
995	Described as "less than 5 cm," or "greater than 4 cm," or "between 4 cm and 5 cm"
	SITE-SPECIFIC CODES WHERE NEEDED
999	Unknown; size not stated Not documented in patient record

Examples:

Mammogram shows 2.5 cm breast malignancy	<i>Code as 025 (2.5 cm = 25 millimeters)</i>
CT of chest shows 4 cm mass in RUL	<i>Code as 040 (4 cm = 40 mm)</i>
Thyroidectomy specimen yields 8 mm carcinoma	<i>Code as 008</i>
Prostate TURP shows 0.6 mm carcinoma	<i>Code as 001 (round up six-tenths of mm)</i>

For schemas that do not use tumor size:

Code	Description
888	Not applicable

Instructions for Coding

1. Refer to general guidelines for Collaborative Staging for timing rules for data collection.
2. Refer to site/histology-specific instructions for additional information. Site/histology-specific instructions replace or over-ride general instructions. Where there are no site/histology-specific instructions, these general instructions apply.
3. Record tumor size information in the following order:
 - a. Record tumor size from the pathology report, if it is available, when the patient receives no radiation or systemic treatment prior to surgery.

Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items

CS Tumor Size, continued

- Example:* Chest x-ray shows 3.5 cm mass; the pathology report from the surgery states that the same mass is malignant and measures 2.8 cm. *Record tumor size as 028.*
- b. If the patient receives preoperative (neoadjuvant) systemic therapy (chemotherapy, hormone therapy, immunotherapy) or radiation therapy, code the largest size of tumor prior to treatment.
- Example:* Patient has a 2.2 cm mass in the oropharynx; fine needle aspiration of mass confirms squamous cell carcinoma. Patient receives course of neoadjuvant combination chemotherapy. Pathologic size of tumor after total resection is 0.8 cm. *Record tumor size as 022.*
- c. Information on size from imaging/radiographic techniques can be used to code size when there is no more specific size information from a pathology or operative report, but it should be taken as low priority, just above a physical exam.
- d. If there is a difference in reported tumor size among imaging and radiographic techniques, record the largest size of tumor reported in the record.
- e. In the infrequent event that the tumor does not respond to neoadjuvant treatment and is, in fact, more extensive after preoperative treatment as determined by the operative or pathology report, code the farthest extension and code CS Tumor Size/Ext Eval as 6, based on pathology/operative report after treatment.
4. Record the exact size of the primary tumor for all sites/histologies except those for which it is stated to be not applicable. If no size is given, code as 999.
- a. Always code the size of the primary tumor, not the size of the polyp, ulcer, cyst, or distant metastasis. However, if the tumor is described as a “cystic mass,” and only the size of the entire mass is given, code the size of the entire mass, since the cysts are part of the tumor itself.
- b. Record the largest dimension or diameter of tumor, whether it is from an excisional biopsy specimen or the complete resection of the primary tumor.
- Example* A 3.3 cm tumor would be 33 millimeters and would be coded as 033.
- Example* Tumor is described as 2.4 x 5.1 x 1.8 cm in size. *Record tumor size as 051.*
- c. Record the size of the invasive component, if given.
- d. If both an *in situ* and an invasive component are present and the invasive component is measured, record the size of the invasive component even if it is smaller.
- Example* Tumor is mixed *in situ* and invasive adenocarcinoma, total 3.7 cm in size, of which 1.4 cm is invasive. *Record tumor size as 014.*
- e. *Additional rule for breast primaries:* If the size of the invasive component is **not** given, record the size of the entire tumor from the surgical report, pathology report, radiology report or clinical examination.
- Example* Infiltrating duct carcinoma with extensive *in situ* component; total size 2.3 cm. *Record tumor size as 023.*
- Example* Duct carcinoma *in situ* covering a 1.9 cm area with focal areas of invasive ductal carcinoma. *Record tumor size as 019.*
- Note:* For breast cancer, document how the size of the tumor was determined in Site Specific Factor field 6. Information from the pathology report can be used to identify *in situ* versus invasive tumor even if exact size is not given. If tumor size is a clinical measurement only in the range 001-989, Site Specific Factor 6 must be coded as 888.
- f. For purely *in situ* lesions, code the size as stated.
- g. Microscopic residual tumor does not affect overall tumor size.
- h. Do **not** add pieces or chips together to create a whole; they may not be from the same location, or they may represent only a very small portion of a large tumor. However, if the pathologist states an aggregate or composite size (determined by fitting the tumor pieces together and measuring the total size), record that size.
- i. If an excisional biopsy is performed and residual tumor at time of resection of the primary is found to be larger than the excisional biopsy, code the size of the residual tumor.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Tumor Size, continued

- j. For an incisional needle biopsy, code tumor size as 999 in the absence of a clinical size. Do not code the tumor size from a needle biopsy unless no residual tumor is found on further resection.
 - k. Record tumor size (lateral dimension) for malignant melanoma. Depth of invasion is coded in a site-specific factor.
 - l. If the tumor is multi-focal or there are multiple tumors being reported as a single primary, code the size of the largest tumor.
5. Special codes
- a. Tumor dimension is to be recorded for all schemas, except as noted below. Other information collected in this field in previous staging systems, such as depth of invasion for melanoma, has been moved to Site-Specific Factors for those sites/histologies.
 - b. If size is not reported, code as 999, which means unknown size or not documented in the patient record.
 - c. The descriptions in code 998 take precedence over any mention of size. Code 998 is used only for the following sites:
 - Esophagus (C15.0-C15.5, C15.8-C15.9): Entire circumference
 - Stomach (C16.0-C16.6, C16.8-C16.9): Diffuse, widespread— $\frac{3}{4}$ or more, linitis plastica
 - Colorectal (M-8220/8221 with /2 or /3): Familial/multiple polyposis
 - Lung and main stem bronchus (C34.0-C34.3, C34.8-C34.9): Diffuse, entire lobe or lung
 - Breast (C50.0-C50.6, C50.8-C50.9): Diffuse
 - d. Code 990, Microscopic focus or foci only; no size is given, should be used when no gross tumor is seen and tumor is only identified microscopically.

Note: the terms microscopic focus, microfocus, and microinvasion are NOT the same as [macroscopic] focal or focus. A macroscopic focus or foci indicates a very small or isolated area, pinpoint, or spot of tumor that may be visible grossly. Only tumor identified microscopically should be coded to 990.

Example Ovary specimen: extensive cystic disease with focal areas of tumor seeding.
Disregard "focal" and code tumor size to 999 unknown.

Example Cervix conization: severe dysplasia with focal areas of microinvasion.
Code tumor size as 990 microscopic focus, no size given.
 - e. Codes 991 through 995 are non-specific size descriptions that, for some sites, could still be used to determine a T category. However, if a specific size is given, the more precise size should be coded in the range 001-989.
 - f. Other special codes in the range 996 to 997 are used on a site-specific basis. See the individual site/histology schemas for further information and definitions.
 - g. **Note:** For the following diagnoses and/or primary sites, size is not applicable. Record as code 888.
 - Disseminated Langerhans cell histiocytosis (Letterer-Siwe disease)
 - Hematopoietic neoplasms
 - Immunoproliferative diseases
 - Leukemia
 - Malignant lymphoma (Hodgkin lymphoma and non-Hodgkin lymphoma)
 - Mast cell tumors
 - Multiple myeloma and other plasma cell tumors
 - Myelodysplastic syndromes
 - Myeloproliferative diseases
 - h. The source of the tumor size (radiographs, endoscopy, pathology specimen, etc.) is documented in the CS Tumor Size/Ext Eval field.
6. It is strongly recommended that the choice of tumor size codes be documented in a related text field on the abstract.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS EXTENSION

Item Length: 2

NAACCR Item #2810

Description

Identifies contiguous growth (extension) of the primary tumor within the organ of origin or its direct extension into neighboring organs. For certain sites such as ovary, discontinuous metastasis is coded in the CS Extension field. See site-specific schemas for detailed codes and coding instructions.

Code	Description	TNM Mapping	SS77 Mapping	SS2000 Mapping
00	In situ; non-invasive	Tis	IS	IS
	SITE/HISTOLOGY-SPECIFIC CODES			
80	Further contiguous extension			
95	No evidence of primary tumor	T0	U	U
99	Unknown extension; primary tumor cannot be assessed; not stated in patient record	TX	U	U

Instructions for Coding

1. Code the farthest documented extension of the primary tumor. Do not include discontinuous metastases to distant sites (these are coded in CS Mets at Dx) except for ovary and corpus uteri (see 2f below).

2. Record extension information in the following order:
 - a. Record extension from the pathology report, if it is available, when the patient receives no radiation or systemic treatment prior to surgery.
 - b. If the patient receives preoperative (neoadjuvant) systemic therapy (chemotherapy, hormone therapy, immunotherapy) or radiation therapy, code the farthest extension identified prior to treatment (clinically).

Example Patient has rectal mass firmly attached to pelvic wall (clinically T4, extension code 60). Patient undergoes preoperative radiation therapy. The pathology report from the low anterior resection shows residual tumor outside the rectum in perimuscular tissue (pathologically T3, extension code 40). *Code extension as 60, because the preoperative treatment apparently “shrank” the tumor away from the pelvic wall.*
 - c. In the infrequent event that the tumor does not respond to neoadjuvant treatment and is, in fact, more extensive after preoperative treatment as determined by the operative or pathology report, code the farthest extension and code CS Tumor Size/Ext Eval as 6, based on pathology/operative report after treatment.

Example Patient found to have an obstructing central lung tumor very close to the main stem bronchus (clinically T2, extension code 20). Patient undergoes six weeks of intensive chemotherapy. At thoracotomy, tumor was observed directly extending into trachea (pathologically T4, extension code 70). *Code extension as 70, because the tumor was noted to be more extensive after the preoperative treatment.*

Example Patient has a 5.5 cm hard, moveable mass in the right breast (clinically T3, extension code 10) and receives preoperative chemotherapy. The pathology report from the modified radical mastectomy shows residual 2.8 cm mass with infiltration of the deep subcutaneous tissues over the mass (pathologically T2, extension code 20). *Code extension as 20, because although the chemotherapy “shrank” the tumor, the residual tumor was found to be more extensive than the clinical presentation.*

Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items

CS Extension, continued

- d. Information on extent of disease from imaging/radiographic techniques can be used to code extension when there is no more specific extension information from a pathology or operative report, but it should be taken as low priority, just above a physical exam.
- e. If an involved organ or tissue is not mentioned in the schema, approximate the location and code it with listed organs or tissues in the same anatomic area.
- f. With the exception of corpus uteri and ovary, all codes represent contiguous (direct) extension of tumor from the site of origin to the organ/structure/tissue represented in the code.

Example Carcinoma of the prostate with extension to pubic bone would be coded 60.
Carcinoma of the prostate with metastases to thoracic spine would be coded in CS Extension to the appropriate code for tumor extension and the metastases to the thoracic spine would be coded in the CS Mets at Dx field.

- 3. Refer to general guidelines for Collaborative Staging for timing rules for data collection.
- 4. Refer to the ambiguous terminology section for terms that constitute tumor involvement or extension.
- 5. If the information in the medical record is ambiguous or incomplete regarding the extent to which the tumor has spread, the extent of disease may be inferred from the T category stated by the physician.
- 6. If the only indication of extension in the record is the physician's statement of a T category from the TNM staging system or a stage from a site-specific staging system, such as Dukes' C, record the numerically lowest equivalent extension code for that T category.
- 7. Some site or histology schemas include designations such as T1, NOS; T2, NOS; Localized, NOS; and other non-specific categories. The NOS is added when there is further breakdown of the category into subsets (such as T1a, T1b, T1c), but the correct subset cannot be determined. The NOS designation, which can appear in both the descriptions of codes and the mapping, is not official AJCC descriptive terminology. The NOS should be disregarded in reports and analyses when it is not a useful distinction. The data collector should only code to a category such as "Stated as T1 NOS" when the appropriate subset (e.g., T1a or T1b) cannot be determined.
- 8. Distant metastases must be coded in the CS Mets at Dx field.
- 9. Do not code CS Extension as in situ if there is any evidence of nodal or metastatic involvement; use the code for Localized, NOS, if there is no better information.
Example Excisional biopsy of breast tumor shows extensive DCIS. Sentinel node biopsy reveals one positive axillary node. *Code CS Extension as 10, localized, NOS, because an in situ tumor theoretically cannot metastasize and apparently an area of invasion was missed by the pathologist.*
- 10. The presence of microscopic residual disease or positive tumor margins does not increase the extension code.
- 11. It is strongly recommended that the choice of extension codes be documented in a related text field on the abstract.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS TUMOR SIZE/EXT EVAL

Item Length: 1

NAACCR Item #2820

Description

Records how the codes for the two items “CS Tumor Size” and “CS Extension” were determined, based on the diagnostic methods employed.

Note: This field is used primarily to describe whether the staging basis for the T category in the TNM system is clinical or pathological.

Code	Description	Staging Basis
0	No surgical resection done. Evaluation based on physical examination, imaging examination, or other non-invasive clinical evidence. No autopsy evidence used.	c
1	No surgical resection done. Evaluation based on endoscopic examination, diagnostic biopsy, including fine needle aspiration biopsy, or other invasive techniques, including surgical observation without biopsy. No autopsy evidence used. <i>Does not meet criteria for AJCC pathologic staging.</i>	c*
2	No surgical resection done, but evidence derived from autopsy (tumor was suspected or diagnosed prior to autopsy)	p
3	Surgical resection performed WITHOUT pre-surgical systemic treatment or radiation OR surgical resection performed, unknown if pre-surgical systemic treatment or radiation performed Evaluation based on evidence acquired before treatment, supplemented or modified by the additional evidence acquired during and from surgery, particularly from pathologic examination of the resected specimen <i>Meets criteria for AJCC pathologic staging.</i>	p
5	Surgical resection performed WITH pre-surgical systemic treatment or radiation; tumor size/extension based on clinical evidence	C
6	Surgical resection performed WITH pre-surgical systemic treatment or radiation, BUT tumor size/extension based on pathologic evidence	Y
8	Evidence from autopsy only (tumor was unsuspected or undiagnosed prior to autopsy)	A
9	Unknown if surgical resection done Not assessed; cannot be assessed Unknown if assessed Not documented in patient record <i>For sites with no TNM schema: not applicable</i>	C

* For some primary sites, code 1 may be a pathologic staging basis, as determined by the site-specific chapter in the *AJCC Cancer Staging Manual, sixth edition*.

Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items

CS Tumor Size/Ext Eval, continued

Instructions for Coding

1. Select the CS Tumor Size/Ext Eval code that documents the report or procedure from which the information about the farthest extension or size of the primary tumor was obtained; this may not be the numerically highest Eval code.
 - Example* Fine needle aspiration biopsy (Eval code 1) confirms adenocarcinoma of prostate. CT scan of pelvis (Eval code 0) shows tumor extension through the prostatic capsule into adjacent connective tissues. *Code CS Tumor Size/Ext Eval as 0 because the CT scan showed more extensive tumor than the biopsy.*
2. For primary sites/histologies where tumor size is not a factor in determining the T category in TNM (see Table 5 in the General Instructions), code CS Tumor Size/Ext Eval on the basis of the CS extension field only.
3. For primary sites where both tumor size and extension determine the T category in TNM (see Table 4 in the General Instructions), select the code that best explains how the information in the CS Tumor Size and CS Extension fields were determined.
 - a. If there is a difference between the derived category for the tumor size and the CS extension, select the evaluation code that reflects how the worse or higher category was determined.
 - Example* Tumor size for a breast cancer biopsy is 020 (maps to T1). There is ulceration of the skin (extension code 51, maps to T4).
Code CS Tumor Size/Ext Eval field as 0, physical examination, because the ulceration information from the physical examination results in a higher T category.
 - b. If the patient had no surgery, use code 0, 1, or 9.
 - Example* Patient has a chest x-ray showing an isolated 4 cm tumor in the right upper lobe. Patient opts for radiation therapy.
Code this field as 0. Staging algorithm would identify information as clinical (c).
 - Example* Colon cancer with colonoscopy and biopsy confirming cancer.
Code this field as 1. Staging algorithm would identify information as clinical (c). The biopsy does not meet the criteria for pathologic staging.
 - Example* Endoscopies for cervix or bladder would be coded as 1 in this field and the staging algorithm would identify the information as clinical (c).
 - Exception* Lung cancer with mediastinoscopy showing direct extension into mediastinum. *Code this field as 1. Staging algorithm would identify information as pathologic (p), because mediastinoscopy is defined as a pathologic procedure in TNM.*
 - c. If the patient had surgery followed by other treatment(s), use code 3 or 9.
 - d. If the size or extension of the tumor determined prior to treatment was the basis for neoadjuvant therapy, use code 5.
 - e. If the size or extension of the tumor was greater after presurgical treatment than before treatment, use code 6. This code is likely to be used infrequently and maps to the “y” intercurrent treatment staging basis.
 - f. If the patient had an autopsy, use code 2 if the diagnosis was known or suspected prior to death. Use code 8 if the malignancy was not known or suspected prior to death.
4. For sites and histologies for which no TNM schema has been defined, such as brain or Kaposi sarcoma, this field is always coded 9, Not Applicable. (See Table 6 in the General Instructions.) For any sites and histologies not listed in Table 6, code to the value that best reflects the diagnostic methods used, whether or not a stage is actually calculated for an individual case. In other words, do not use code 9 when a case has a histology that is excluded from staging but the site does have a TNM schema defined, for example, a sarcoma of the breast. In those cases, use code 9 only when the nature of the diagnostic methods is actually unknown.

Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items

CS Tumor Size/Ext Eval, continued

5. Code 0 includes imaging studies such as standard radiography, special radiographic projections, tomography, computerized tomography (CT), ultrasonography (US), lymphography, angiography, scintigraphy (nuclear scans), magnetic resonance imaging (MRI), positron emission tomography (PET) scans, spiral scanning (CT or MRI) and other non-invasive methods of examining tissues.
6. Codes 0-3 are oriented to the AJCC staging basis. In general, Code 1 includes microscopic analysis of tissue that is insufficient to meet the requirements for pathologic staging in the TNM system. However, pathologic staging requirements vary by site; for some site schemas, code 1 may be classified as pathologic. For specific classification rules, refer to the *AJCC Cancer Staging Manual, sixth edition*. For example, a total cystectomy is required to pathologically stage a bladder cancer. Any tissue removed during another procedure, such as a transurethral resection of a bladder tumor, would not meet the requirements for pathologic staging and should be coded to 1 in this field. Code 1 also includes observations at surgery, such as an exploratory laparotomy in which unresectable pancreatic cancer is identified, where further tumor extension is not biopsied.
7. For most schemas, Code 3 is considered pathologic staging. For most schemas, use code 3 for a biopsy of tumor extension that meets the requirements for pathologic staging basis. In other words, according to TNM rules, if the biopsy documents the highest T category, the biopsy meets the requirements for pathologic staging basis and the CS Tumor Size/Ext Eval field should be coded to 3. For example, if a prostate cancer patient has a biopsy of the rectum that shows microscopic involvement of the rectal wall (T4), according to the *AJCC Cancer Staging Manual sixth edition* that patient meets the requirements for pathologic staging in the T category.
8. The Eval fields should be coded based on how the information was obtained, even if the information in the related field (Tumor Size, Regional Nodes, or Mets at Dx) is unknown. For example, even if it is not possible to determine the tumor size or extension and the Extension field is coded as 99, the registrar still knows what procedures were used to try to determine those fields. In other words, just because the tumor size is coded 999, the Eval field does not have to be coded 9.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS LYMPH NODES

Item Length: 2

NAACCR Item #2830

Description

Identifies the regional lymph nodes involved with cancer at the time of diagnosis.

Code	Description	TNM Mapping	SS77 Mapping	SS2000 Mapping
00	None; no regional lymph node involvement	N0	None	None
	SITE/HISTOLOGY-SPECIFIC CODES			
99	Unknown; regional lymph nodes cannot be assessed; not stated in patient record	NX	U	U

For schemas that do not use the CS Lymph Nodes field:

Code	Description
88	Not applicable

Instructions for Coding

1. Record the specific regional lymph node chain farthest from the primary site that is involved by tumor either clinically or pathologically.
 - a. Regional lymph nodes are listed for each site/histology. In general, the regional lymph nodes in the chain(s) closest to the primary site have the lower codes. Nodes farther away from the primary or in farther lymph node chains have higher codes. Record the highest applicable code.

Exception The higher codes for 'Regional lymph nodes, NOS'; 'Lymph nodes, NOS'; 'Stated as N1, no other information'; 'Stated a N2a, no other information', and so forth, should only be used when there is no available information as to the name(s) of the regional nodes involved.

Example Peribronchial lymph nodes are positive on fine needle aspiration biopsy. Contralateral mediastinal mass noted on CT scan but not biopsied. Patient chooses radiation therapy as primary treatment.
Use the code for contralateral mediastinal lymph node involvement as it is higher than the code for peribronchial lymph nodes.
 - b. Record involved regional lymph nodes from the pathology report, if it is available, when the patient receives no radiation or systemic treatment prior to surgery.
 - c. If there is a discrepancy between clinical information and pathologic information about the same lymph nodes, the pathologic information takes precedence if no preoperative treatment was administered.

Example Axillary lymphadenopathy stated as "suspicious for involvement" noted on physical exam. After axillary dissection, all lymph nodes are negative.
Code CS Lymph Nodes as 0, no regional lymph node involvement.
 - d. *For inaccessible sites*, record CS Lymph Nodes as Code 00 (None) rather than Code 99 (Unknown) when the clinician proceeds with standard treatment of the primary site for clinically localized or early (for example, T1, T2) stage disease, since this action presumes that there are no involved regional lymph nodes that would otherwise alter the treatment approach. Code 99 can and should be used in situations where there is reasonable doubt that the tumor is no longer localized and there is no documentation of involved regional lymph nodes.

Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items

CS Lymph Nodes, continued

- e. If there is direct extension of the primary tumor into a regional lymph node, record the involved node in this field.
- f. If the patient receives preoperative (neoadjuvant) systemic therapy (chemotherapy, hormone therapy, immunotherapy) or radiation therapy, code the farthest involved regional lymph nodes, based on information prior to surgery.

Example Patient has a hard matted mass in the axilla (code 50) and a needle biopsy of the breast that confirms ductal carcinoma. Patient receives three months of chemotherapy. The pathology report from the modified radical mastectomy shows only scar tissue in the axilla with no involvement of axillary lymph nodes (Negative, code 00). *Code CS Lymph Nodes as 50 because the chemotherapy apparently "sterilized" the lymph nodes.*

- g. In the infrequent event that clinically involved regional lymph nodes do not respond to neoadjuvant treatment and are, in fact, more extensively involved after preoperative treatment as determined by the operative or pathology report, code the farthest extension and code CS Reg Nodes Eval as 6, based on pathology/operative report after treatment.

Example Patient has needle biopsy-proven prostate cancer with no mention of involved lymph nodes on physical examination (Negative, code 00). He receives Lupron while deciding whether to undergo a radical prostatectomy. At the time of surgery, a laparoscopic pelvic node biopsy is reported to show metastases (Regional nodes involved, code 10) to lymph nodes and the prostatectomy is canceled. *Code CS Lymph Nodes as 10 because the preoperative treatment (Lupron) had no effect on the lymph nodes.*

- 2. Use code 00 for lymph node involvement when the CS Extension is coded in situ, even if no lymph nodes are removed, since "in situ" by definition means noninvasive. If there is evidence of nodal involvement associated with a tumor described as in situ, it would indicate that an area of invasion was missed and the primary tumor is not an in situ lesion, so lymph nodes can be coded as appropriate for the case.
- 3. For solid tumors, the terms "fixed" or "matted" and "mass in the hilum, mediastinum, retroperitoneum, and/or mesentery" (with no specific information as to tissue involved) are considered involvement of lymph nodes.
 - a. Any other terms, such as "palpable," "enlarged," "visible swelling," "shotty," or "lymph-adenopathy" should be ignored, unless there is a statement of involvement by the clinician.

Exception The terms *adenopathy*, *enlargement*, and *mass in the hilum or mediastinum* should be coded as involvement for lung primaries only.
 - b. For lymphomas, *any* positive mention of lymph nodes indicates involvement of those lymph nodes.
 - c. Regional lymph nodes are not palpable for inaccessible sites such as bladder, kidney, prostate, esophagus, stomach, lung, liver, corpus uteri and ovary. The best description concerning regional lymph nodes will be on imaging studies or in the surgeon's evaluation at the time of exploratory surgery or definitive surgery. If regional lymph nodes for these inaccessible sites are not mentioned on imaging or exploratory surgery, they are presumed to be clinically negative.
 - d. The terms "homolateral," "ipsilateral," and "same side" are used interchangeably.
 - e. Any unidentified nodes included with the resected primary site specimen are to be coded as regional lymph nodes, NOS.
 - f. Where more specific categories are provided, the codes for "regional lymph node(s), NOS"; "lymph nodes, NOS"; and "Stated as N_, no additional information" should be used *only* after an exhaustive search for more specific information.

Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items

CS Lymph Nodes, continued

4. When size of involved regional lymph nodes is required, code from pathology report, if available.
 - a. Code the size of the metastasis, not the entire node, unless otherwise stated in site-specific schemas. The size of the metastasis within the lymph node can be inferred if the size for the entire node falls within one of the codes; for example, a single involved node 1.5 cm in size can be coded to “single lymph node \leq 2 cm” because the metastasis cannot be larger than 1.5 cm.
5. If the only indication of lymph node involvement in the record is the physician’s statement of an N category from the TNM staging system or a stage from a site-specific staging system, such as Dukes’ C, record the numerically lowest equivalent CS Lymph Nodes code for that N category.
 - a. If there is a discrepancy between documentation in the medical record and the physician’s assignment of TNM, the documentation takes precedence. Cases of this type should be discussed with the physician who assigned the TNM.
 - b. If the information in the medical record is ambiguous or incomplete regarding the extent to which the tumor has spread, lymph node involvement may be inferred from the N category stated by the physician.
6. Some site or histology schemas include designations such as N1, NOS; N2, NOS, and other non-specific categories. The NOS is added when there is further breakdown of the category into subsets (such as N1a, N1b, N1c), but the correct subset cannot be determined. The NOS designation, which can appear in both the descriptions of codes and the mapping, is not official AJCC descriptive terminology. The NOS should be disregarded in reports and analyses when it is not a useful distinction. The data collector should only code to a category such as “Stated as N1 NOS” when the appropriate subset (e.g., N1a or N1b) cannot be determined.
7. For colon, rectosigmoid and rectum primaries, if there is a statement about tumor nodule(s) in the pericolic or perirectal fat, use the following guidelines for coding regional lymph node involvement:
Code as regional lymph node involvement if the nodule has a smooth contour.
Code as tumor extension if the nodule has an irregular contour.
8. It is strongly recommended that the choice of regional lymph node codes be documented in a related text field on the abstract.

CODING REGIONAL LYMPH NODES FOR HEAD AND NECK SITES

For head and neck sites, regional lymph node information is coded in several fields. The CS Lymph Nodes field contains information about the nodes involved, their number and laterality. Site-Specific Factors 1 and 2 are used to code the size of involved lymph nodes and the presence of extracapsular extension. Site-Specific Factors 3 through 6 are used to code the presence or absence of lymph node involvement in each of 7 different levels and other groups defined by AJCC. The definitions of the levels are the same for all applicable head and neck sites. One digit is used to represent lymph nodes of a single level, with the three digits of Site-Specific Factor 3 representing lymph nodes of, respectively, Levels I-III; the digits of Site-Specific Factor 4 representing lymph nodes of Levels IV and V and the retropharyngeal nodes; the digits of Site-Specific Factor 5 representing lymph nodes of Levels VI and VII and the facial nodes; and the digits of Site-Specific Factor 6 representing the remaining Other groups as defined by AJCC. In each digit, a code 1 means Yes, the nodes are involved. See Figure 2a for the layout of Site-Specific Factors 3 through 6 and Figure 2b for the interpretation of a coded example.

Figure 2a. Layout of Site-Specific Factors for Head and Neck Sites

SSF 3	Levels I-III	<u> </u> I	<u> </u> II	<u> </u> III
SSF 4	Levels IV-V, retropharyngeal (RP)	<u> </u> IV	<u> </u> V	<u> </u> RP
SSF 5	Levels VI-VII, Facial (F)	<u> </u> VI	<u> </u> VII	<u> </u> F
SSF 6	Other groups Parapharyngeal (PP), Parotid (PA), Suboccipital (S)	<u> </u> PP	<u> </u> PA	<u> </u> S

Figure 2b. Example and Interpretation of Site-Specific Factors for Head and Neck Sites

Example: Left Radical Neck Dissection: 2 positive parotid node (< 3 cm with extra-capsular extension), 1 positive buccal (facial) node (2 cm), and 1 positive 2 cm submandibular node.

SSF 3	Levels I-III	<u> 1 </u> I	<u> 0 </u> II	<u> 0 </u> III
SSF 4	Levels IV-V, retropharyngeal (RP)	<u> 0 </u> IV	<u> 0 </u> V	<u> 0 </u> RP
SSF 5	Levels VI-VII, Facial (F)	<u> 0 </u> VI	<u> 0 </u> VII	<u> 1 </u> F
SSF 6	Other groups Parapharyngeal (PP), Parotid (PA), Suboccipital (S)	<u> 0 </u> PP	<u> 1 </u> PA	<u> 0 </u> S

<u>Stored in database as</u>		<u>Interpretation</u>
SSF 3	100	Level 1 only
SSF 4	000	All nodes neg
SSF 5	001	Facial nodes only
SSF 6	010	Parotid nodes only

CS Lymph Nodes, continued

Unknown

In Site-Specific Factors 3-6 for lymph node levels, use code 9 only when it is unknown if lymph nodes are involved. Within each of the Site-Specific Factors 3-6, do not code 9 in some positions and 0 or 1 in other positions. If specific information is available about the positive or negative status of some but not all nodes in any one level or group, assume that the rest of the nodes in the same Site-Specific Factor are negative and code accordingly.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

NOS

When the only information available is “Regional nodes, NOS” or “Cervical nodes, NOS” or “Internal jugular lymph nodes, NOS” or “Lymph nodes, NOS,” code 0 in all digits of Site-Specific Factors 3-6.

Example 1 A carcinoma of the base of tongue involves bilateral submandibular nodes and left upper, mid-, and lower jugular nodes, the largest measuring 4 cm. There is no extracapsular extension. These are level I, II, III, and IV lymph nodes according to AJCC definitions. CS Lymph Nodes is coded 40 (bilateral or contralateral nodes). Site-Specific Factor 1 is coded 040 indicating the largest size. Site-Specific Factor 2 is coded 000 for no extracapsular extension. Site-Specific Factor 3 is coded 111, to show that levels I, II, and III are involved. Site-Specific Factor 4 is coded 100 to show that level IV is involved. Site-Specific Factors 5 and 6 are each coded 000, since no other nodes are involved.

Example 2 Laryngeal biopsy with squamous cell carcinoma, no other information available. CS Lymph Nodes is coded 99. Site-Specific factors 1-6 are each coded 999, since no information is available regarding lymph node involvement.

Example 3 Patient diagnosed elsewhere with carcinoma of oropharynx with cervical lymph node involvement. No other information available. CS Lymph Nodes is coded 50 (regional nodes, NOS, not stated if ipsilateral, bilateral, or contralateral, or if single or multiple). Site-specific Factors 1 and 2 are each coded 999. Site-Specific Factors 3-6 are each coded 000.

Definitions of Levels for Head and Neck Sites

The definitions of the levels and the lymph node chains included in each level are as follows:

Level I contains the submental and submandibular triangles bounded by the anterior and posterior bellies of the digastric muscle, and the hyoid bone inferiorly, and the body of the mandible superiorly.

Submandibular	Submaxillary	Submental
---------------	--------------	-----------

Level II contains the upper jugular lymph nodes and extends from the level of the skull base superiorly to the hyoid bone inferiorly.

Jugulodigastric (subdigastric)	Upper deep cervical	Upper jugular
-----------------------------------	---------------------	---------------

Level III contains the middle jugular lymph nodes from the hyoid bone superiorly to the level of the lower border of the cricoid cartilage inferiorly.

Middle deep cervical	Mid-jugular
----------------------	-------------

Level IV contains the lower jugular lymph nodes from the level of the cricoid cartilage superiorly to the clavicle inferiorly.

Jugulo-omohyoid (supraomohyoid)	Lower deep cervical	Lower jugular
------------------------------------	---------------------	---------------

Level V contains the lymph nodes in the posterior triangle bounded by the anterior border of the trapezius muscle posteriorly, the posterior border of the sternocleidomastoid muscle anteriorly, and the clavicle inferiorly. For descriptive purposes, Level V may be further subdivided into upper, middle, and lower levels corresponding to the superior and inferior planes that define Levels II, III, and IV.

Posterior cervical
Posterior triangle (spinal accessory and transverse cervical) (upper, middle, and lower, corresponding to the levels that define upper, middle, and lower jugular nodes)

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

Level VI contains the lymph nodes of the anterior central compartment from the hyoid bone superiorly to the suprasternal notch inferiorly. On each side, the lateral boundary is formed by the medial border of the carotid sheath.

Anterior deep cervical	Paratracheal	Pretracheal
Laterotracheal	Prelaryngeal (Delphian)	Recurrent laryngeal
Paralaryngeal		

Level VII contains the lymph nodes inferior to the suprasternal notch in the superior mediastinum.

Upper mediastinal

Other groups

Buccinator (facial)	Periparotid and	Retropharyngeal
Nasolabial	intraparotid	Sub-occipital
Parapharyngeal	Preauricular	

CODING REGIONAL LYMPH NODES FOR BREAST

Coding regional lymph node involvement for breast cancers is more complex than for many other sites, especially when dealing with isolated tumor cells (ITCs) and micrometastases. The following may help clarify the reasons behind the codes in CS Lymph Nodes and Site-Specific Factors 3-5. For a more detailed explanation, see the section in the breast chapter of the *AJCC Cancer Staging Manual, 6th edition*, called "Considerations for Evidence-Based Changes to the *AJCC Cancer Staging Manual, 6th edition*," beginning on page 229.

Isolated Tumor Cells (ITCs). Pathologists can detect isolated tumor cells (ITCs) spread from a breast cancer into regional lymph nodes. These are very small deposits of tumor cells, so small that they are *not* considered significant for assigning stage. They usually do not show evidence of malignant activity in the nodes, such as proliferation or stromal reaction. To be considered ITCs, they must be single tumor cells or small clusters not more than 0.2 mm. As more data are collected about these ITCs, their prognostic significance may be better understood. **At this time, nodes with only these ITCs are NOT considered positive nodes.** These ITCs are most often found using immunohistochemistry tests on sentinel lymph node specimens. The ITCs may sometimes also be seen on routine H&E-stained sections.

Hematoxylin and Eosin (H & E). (from "Hematoxylin & Eosin": (The Routine Stain)), by H. Skip Brown, BA, HT(ASCP), from: <http://www.sigmaldrich.com/img/assets/7361/Primer-H&Emay04.pdf>

In histology, the standard or routine stain is the hematoxylin and eosin stain, better known as the "H&E" stain. With rare exceptions, every specimen being examined will first receive an H&E stain to give the laboratorian a visible look at the nucleus of the cells and their present state of activity. With most disease states there is abnormal growth and/or division in the nucleus of the cells. The hematoxylin and eosin stain uses two separate dyes, one staining the nucleus and the other staining the cytoplasm and connective tissue. Hematoxylin is a dark purplish dye that will stain the chromatin (nuclear material) within the nucleus, leaving it a deep purplish-blue color. Eosin is an orangish-pink to red dye that stains the cytoplasmic material including connective tissue and collagen, and leaves an orange-pink counterstain. This counterstain acts as a sharp contrast to the purplish-blue nuclear stain of the nucleus, and helps identify other entities in the tissues such as cell membrane (border), red blood cells, and fluid.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

Immunohistochemistry (IHC). Immunohistochemistry (IHC) tests use antibodies to stain for proteins of interest in tissue specimens. The IHC test for metastatic breast cancer in lymph nodes uses antibodies to cytokeratin, so the test may be called “cytokeratin staining.” Other IHC tests are used on the primary breast tumor, rather than the lymph nodes, to assess estrogen and progesterone receptors and HER-2 neu (human epidermal growth factor receptor). In SSF 4, code only IHC results for ITCs in LYMPH NODES.

Molecular Study: Reverse Transcriptase/Polymerase Chain Reaction (RT-PCR). An even more sensitive test used to detect ITCs in lymph nodes is RT-PCR, a molecular test looking for expression of genes of interest. This test is rarely done.

Micrometastasis. When the tumor deposits in the lymph nodes are larger than 0.2 mm but not larger than 2.0 mm, they are defined as micrometastasis. **Nodes with micrometastasis ARE considered positive for staging.**

In coding CS Lymph Nodes and Site-Specific Factors 3-5, the important things to abstract are the size of the tumor detected in the lymph nodes and the methods of detection. The table below may help in coding this information. Note that the table includes codes for axillary nodes only, not internal mammary nodes. The table is followed by examples to illustrate likely coding situations.

To use the table, identify the group (numbered I-VI) of applicable rows based on the information in column 2 that best represents the information in the case. Within that group, find the row or rows that represent the information in the case, and read right to the last four columns to find the codes to use. The group numbers are for convenience in using this chart only, and do not correlate with any anatomic groups of nodes.

GIVEN THIS INFORMATION...				USE THESE CODES...			
		Row Number	IHC and/or Mol Studies Done, or Method of Detection/ Verification	CS Lymph Nodes	SSF3 (Number pos axillary nodes)	SSF4 (IHC)	SSF5 (Mol)
I.	Clinical information only; no pathological information used to code CS Lymph Nodes; no nodes examined pathologically, nodes clinically NEGATIVE	1.	None; does not apply	00	098	000	000
II.	Clinical information only; no pathological information used to code CS Lymph Nodes; no nodes examined pathologically, nodes clinically POSITIVE	2.	None; does not apply	50, 60, or 99	098	888	888

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

III.	Nodes examined pathologically, nodes negative; no Isolated Tumor Cells (ITCs) NOTE: SSF 4 and 5 are coded independently of each other.	3.	Immunohistochemistry (IHC) (cytokeratin staining) not done, OR unknown if done	00	000	000	
		4.	IHC done, neg for tumor	00	000	001	
		5.	Molecular studies not done, OR unknown if done	00	000		000
		6.	Molecular studies done, neg for tumor	00	000		001
IV.	Nodes examined pathologically, Isolated Tumor Cells (ITCs) ONLY; Single tumor cells, or clusters \leq 0.2mm OR Immunohistochemistry (IHC) pos, NOS NOTE: SSF 4 and 5 are coded independently of each other.	7.	H&E (routine stained slides)	05	000	888	888
		8.	H&E neg, immunohistochemistry (IHC) (cytokeratin staining) not done, OR unknown if done	00	000	000	
		9.	H&E neg, IHC done, neg for ITCs	00	000	001	
		10.	H&E neg, IHC done, pos for ITCs	00	00	002	
		11.	H&E neg, IHC done, pos but size of deposits not stated	00	000	009	
		12.	H&E neg, molecular studies not done, or unknown if done	00	000		000
		13.	H&E neg, molecular studies done, neg for tumor	00	000		001
		14.	H&E neg, molecular studies done, pos for ITCs	00	000		002
		V.	Nodes examined pathologically Tumor > 0.2mm and \leq 2.0mm (Micrometastasis)	15.	H&E neg, micromets on IHC (cytokeratin staining) ONLY	13	001-097
16.	H&E pos for micromets			15	001-097	888	888
VI.	Nodes examined pathologically Tumor > 2.0mm; positive lymph nodes	17.	Does not apply	25 or higher	001-097	888	888

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

Examples for Each Group

Group I Example

1. Nodes clinically negative, patient refused further workup. [Row number 1]

Group II Examples

1. Fixed and matted ipsilateral axillary nodes clinically, patient had pre-op chemotherapy. Subsequent modified radical mastectomy showed negative axillary nodes. (CS Reg Nodes Eval = 5 in this case.) [Row number 2]
2. Axillary nodes clinically positive, patient refused further workup. [Row number 2]

Group III Examples

1. Sentinel nodes neg on H&E. IHC (cytokeratin stain) performed, negative for ITCs. Molecular studies not done. [Rows 4 and 5]
2. Modified radical mastectomy, path report with 12 lymph nodes neg for tumor, no special stains, cytokeratin, IHC, or molecular studies performed on lymph nodes. [Rows 3 and 5]
3. Sentinel nodes neg on H&E. Unknown if IHC done. RT-PCR done, negative for ITCs. [Rows 3 and 6]

Group IV Examples

1. Sentinel nodes neg on H&E. IHC (cytokeratin stain) performed, positive for ITCs. Unknown if molecular studies done. [Rows 10 and 12]
2. Sentinel nodes initially neg on H&E. IHC performed, positive for ITCs. No molecular studies done. ITCs then verified on H&E slides of the sentinel nodes. [Row 7 ONLY]
3. Class 3 case abstracted from clinical history. Sentinel nodes neg on H&E. IHC on sentinel nodes was positive, NOS. Molecular studies not mentioned. [Rows 11 and 12]
4. Sentinel nodes neg on H&E. Cytokeratin stain showed clusters of tumor cells in the node up to 0.15 mm. RT-PCR was pos for ITCs. [Rows 10 and 14]
5. Sentinel nodes neg on H&E. Unknown if IHC performed. RT-PCR study done, neg for ITCs. [Rows 8 and 13]
6. Sentinel nodes neg on H&E. IHC and RT-PCR negative for tumor. [Rows 9 and 13]

Group V Examples

1. Path report, final diagnosis: "Lymph Nodes: one of three sentinel lymph nodes positive for capsular micrometastases." Microscopic description: "Sections of the first submitted sentinel lymph node demonstrate normal nodal architecture, however, on cytokeratin stain, micrometastases are noted in the capsule." [Row 15]
2. Path report, final diagnosis: "Lymph Nodes: one of three sentinel lymph nodes positive for capsular micrometastases." Microscopic description: "Sections of the first submitted sentinel lymph node demonstrate micrometastases in the capsule." No special studies are mentioned in the report. [Row 16]

Group VI Example

1. Axilla neg on palpation. Modified radical mastectomy, 2/14 nodes positive. Largest metastasis 0.8 cm. [Row 17]

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

Page left blank.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS REG NODES EVAL

Item Length: 1

NAACCR Item #2840

Description

Records how the code for the item "CS Lymph Nodes" was determined, based on the diagnostic methods employed.

Code	Description	Staging Basis
0	No regional lymph nodes removed for examination. Evaluation based on physical examination, imaging, or other non-invasive clinical evidence. No autopsy evidence used.	c
1	No regional lymph nodes removed for examination. Evaluation based on endoscopic examination, diagnostic biopsy including fine needle aspiration of lymph node(s) or other invasive techniques, including surgical observation without biopsy. No autopsy evidence used. <i>Does not meet criteria for AJCC pathologic staging.</i>	c
2	No regional lymph nodes removed for examination, but evidence derived from autopsy (tumor was suspected or diagnosed prior to autopsy).	p
3	Regional lymph nodes removed for examination (removal of at least 1 lymph node) WITHOUT pre-surgical systemic treatment or radiation OR lymph nodes removed for examination, unknown if pre-surgical systemic treatment or radiation performed <i>Meets criteria for AJCC pathologic staging.</i>	p
5	Regional lymph nodes removed for examination WITH pre-surgical systemic treatment or radiation, and lymph node evaluation based on clinical evidence.	c
6	Regional lymph nodes removed for examination WITH pre-surgical systemic treatment or radiation, BUT lymph node evaluation based on pathologic evidence.	y
8	Evidence from autopsy; tumor was unsuspected or undiagnosed prior to autopsy.	a
9	Unknown if lymph nodes removed for examination Not assessed; cannot be assessed Unknown if assessed Not documented in patient record <i>For sites that have no TNM staging:</i> Not applicable	c

Instructions for Coding

1. Select the CS Reg Nodes Eval code that documents the report or procedure from which the information about the farthest involved regional lymph nodes was obtained; this may not be the numerically highest eval code.

Example Modified radical neck dissection for hypopharyngeal cancer shows one lower jugular node involved (CS Reg LN code 10, Eval code 3). Physical exam shows hard, matted scalene (transverse cervical) node presumed to contain metastasis (CS Reg LN code 32, Eval code 0). *Code CS Reg Nodes Eval as 0 since the scalene node involvement was determined clinically rather than by examination of tissue.*

2. For sites and histologies for which no TNM schema has been defined, such as brain or Kaposi sarcoma, this field is always coded 9, Not Applicable. (See Table 6 in the General Instructions.) For

Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items

CS Reg Nodes Eval, continued

any sites and histologies not listed in Table 6, code to the value that best reflects the diagnostic methods used, whether or not a stage is actually calculated for an individual case. In other words, do not use code 9 when a case has a histology that is excluded from staging but the site does have a TNM schema defined, for example, a sarcoma of the breast. In those cases, use code 9 only when the nature of the diagnostic methods is actually unknown.

3. Select the code that best explains how the information in the CS Lymph Nodes field was determined.
 - a. If the patient had no removal of lymph node(s), use code 0, 1, or 9.

Example Prostate cancer with laparoscopic lymph node biopsy showing involved nodes; radical prostatectomy canceled.
Code CS Reg Node Eval as 3. Staging algorithm would identify information as pathologic (p). According to AJCC, a positive biopsy of one or more regional lymph nodes is sufficient to meet the pathologic staging basis for prostate cancer.

Example Lung cancer with CT scan or MRI showing involved contralateral mediastinal nodes.
Code CS Reg Node Eval as 0. Staging algorithm would identify information as clinical (c).
 - b. If the patient had removal of lymph node(s) surgery followed by other treatment(s), use code 3 or 9.
 - c. If the patient receives preoperative (neoadjuvant) systemic therapy (chemotherapy, hormone therapy, immunotherapy) or radiation therapy, the clinical status of lymph nodes takes precedence (code 5).
 - d. If the size, number or extension of regional lymph node involvement determined prior to treatment was the basis for neoadjuvant therapy, use code 5. However, if more extensive tumor is found during lymph node examination after neoadjuvant therapy, use code 6.
 - e. If the patient had an autopsy, use code 2 if the diagnosis was known or suspected prior to death. Use code 8 if the malignancy was not known or suspected prior to death.
4. Code 0 includes imaging studies such as standard radiography, special radiographic projections, tomography, computerized tomography (CT), ultrasonography (US), lymphography, angiography, scintigraphy (nuclear scans), magnetic resonance imaging (MRI), positron emission tomography (PET) scans, spiral scanning (CT or MRI) and other non-invasive methods of examining tissues.
5. Codes 0-3 are oriented to the AJCC staging basis. Code 1 includes microscopic analysis of tissue insufficient to meet the requirements for pathologic staging in the TNM system. For example, a needle biopsy of an axillary lymph node will document that a lymph node is involved by breast cancer, but does not meet the requirement for removal of a sufficient number of lymph nodes so that the highest N stage can be assessed. Pathologic staging requirements vary by site; for some site schemas, code 1 may be classified as pathologic. For specific classification rules, refer to the *AJCC Cancer Staging Manual, sixth edition*. Code 1 also includes observations at surgery, such as abdominal exploration at the time of a colon resection, where regional lymph nodes are not biopsied.
6. Code 3 maps to pathologic staging across all sites. Use code 3 if the lymph node procedure meets the requirements for pathologic staging basis of regional lymph nodes. The requirements vary among sites as to the location and number of lymph nodes involved, the size of the involved nodes, and other characteristics. For prostate cancer, a positive biopsy of a single regional lymph node is sufficient to assign CS Reg Nodes Eval code 3 to the case.
7. The Eval fields should be coded based on how the information was obtained, even if the information in the related field (Tumor Size, Regional Nodes, or Mets at Dx) is unknown. For example, even if it is not possible to determine the tumor size or extension and the Extension field is coded as 99, the registrar still knows what procedures were used to try to determine those fields. In other words, just because the tumor size is coded 999, the Eval field does not have to be coded 9.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

REGIONAL NODES POSITIVE

Item Length: 2

NAACCR Item #820

Description

Records the exact number of regional lymph nodes examined by the pathologist and found to contain metastases.

Code	Description
00	All nodes examined are negative.
01-89	1-89 nodes are positive. (Code exact number of nodes positive)
90	90 or more nodes are positive.
95	Positive aspiration or core biopsy of lymph node(s) was performed.
97	Positive nodes are documented, but the number is unspecified.
98	No nodes were examined.
99	It is unknown whether nodes are positive; not applicable; not stated in patient record.

Instructions for Coding

1. Record information about only regional lymph nodes in this field. Involved distant lymph nodes should be coded in the "CS Mets at Dx" field.
2. Rules for coding Regional Nodes Positive are the same for both in situ and invasive cases.
3. This field is based on pathologic information only. If no lymph nodes were removed for examination, or if a lymph node drainage area was removed but no lymph nodes were found, code as 98.
4. Record the total number of regional lymph nodes removed and found to be positive by pathologic examination.
 - a. The number of regional lymph nodes positive is cumulative from all procedures that removed lymph nodes through the completion of surgeries in the first course of treatment.
 - b. This field is to be recorded regardless of whether the patient received preoperative treatment.
5. Any combination of positive aspirated, biopsied, sampled or dissected lymph nodes should be coded to 97 if the number of involved nodes cannot be determined on the basis of cytology or histology.
6. For the following primary sites and histologies, the Regional Nodes Positive field is always coded as 99.
 - Placenta
 - Brain and Cerebral Meninges
 - Other Parts of Central Nervous System
 - Hodgkin and non-Hodgkin Lymphoma
 - Hematopoietic, Reticuloendothelial, Immunoproliferative and Myeloproliferative Neoplasms
 - Other and Ill-Defined Primary Sites
 - Unknown Primary Site

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

REGIONAL NODES EXAMINED

Item Length: 2

NAACCR Item #830

Description

Records the total number of regional lymph nodes that were removed and examined by the pathologist.

Code	Description
00	No nodes were examined.
01-89	1-89 nodes were examined. (Code the exact number of regional lymph nodes examined.)
90	90 or more nodes were examined.
95	No regional nodes were removed, but aspiration or core biopsy of regional nodes was performed.
96	Regional lymph node removal was documented as a sampling, and the number of nodes is unknown/not stated.
97	Regional lymph node removal was documented as a dissection, and the number of nodes is unknown/not stated.
98	Regional lymph nodes were surgically removed, but the number of lymph nodes is unknown/not stated and not documented as a sampling or dissection; nodes were examined, but the number is unknown.
99	It is unknown whether nodes were examined; not applicable or negative; not stated in patient record.

Instructions for Coding

1. Record information about only regional lymph nodes in this field. Distant lymph node information should be coded in the "CS Mets at Dx" field.
2. Rules for coding Regional Nodes Examined are the same for in situ and invasive cases.
3. This field is based on pathologic information only. If no lymph nodes were removed for examination, or if a lymph node drainage area was removed but no lymph nodes were found, code as 00. If it is unknown whether nodes were removed or examined, code as 99.
4. Record the total number of regional lymph nodes removed and examined by the pathologist.
 - a. The number of regional lymph nodes examined is cumulative from all procedures that removed lymph nodes through the completion of surgeries in the first course of treatment.
 - b. If lymph nodes are aspirated and other lymph nodes are removed, use code 98.
 - c. This field is to be recorded regardless of whether the patient received preoperative treatment.
5. If a lymph node biopsy was performed, code the number of nodes removed, if known. If the number of nodes removed by biopsy is not known, use code 96.
6. For the following primary sites and histologies, the Regional Nodes Examined field is always coded as 99.
 - Brain and Cerebral Meninges
 - Hematopoietic, Reticuloendothelial, Immunoproliferative and Myeloproliferative Neoplasms
 - Hodgkin and non-Hodgkin Lymphoma
 - Other and Ill-Defined Primary Sites
 - Other Parts of Central Nervous System
 - Placenta
 - Unknown Primary Site

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS METS AT DX

Item Length: 2

NAACCR Item #2850

Description

Identifies the distant site(s) of metastatic involvement at time of diagnosis.

Code	Description	TNM Mapping	SS77 Mapping	SS2000 Mapping
00	No; none	M0	None	None
10	Distant lymph node(s)	M1	D	D
40	Distant metastases except code 10 Distant metastasis, NOS Carcinomatosis	M1	D	D
	SITE/HISTOLOGY-SPECIFIC CODES WHERE NEEDED			
50	(40) + (10)	M1	D	D
99	Unknown; distant metastasis cannot be assessed; not stated in patient record	MX	U	U

For schemas that do not use the CS Mets at Dx field:

Code	Description
88	Not applicable

Instructions for Coding

1. This field represents distant metastases (the TNM M component or distant stage in Summary Staging) at the time of diagnosis. In other words, when the patient was diagnosed, tumor had already spread indirectly (through vascular or lymph channels) to a site remote from the primary tumor.

Note: The structure of the CS Mets at Dx field is based on the M category of TNM. In some schemas, there may be additional items in CS Extension or CS Lymph Nodes that map to distant stage in Summary Staging (77 and/or 2000) and there may be some items in CS Mets at Dx that map to regional stage in Summary Staging. Regardless of where such items are recorded, the staging algorithms will properly account for the information.

2. Assign the highest applicable code for metastasis at diagnosis, whether the determination was clinical or pathological and whether or not the patient had any preoperative systemic therapy.
3. Metastasis known to have developed after the extent of disease was established (also referred to as progression of disease) should not be recorded in the CS Mets at Dx field.
4. Record CS Mets at Dx as Code 00 (None) rather than Code 99 (Unknown) when the clinician proceeds with standard treatment of the primary site for clinically localized or early (for example, T1, T2) stage disease, since this action presumes that there are no distant metastasis that would otherwise alter the treatment approach. Code 99 can and should be used in situations where there is reasonable doubt that the tumor is no longer localized and there is no documentation of distant metastases.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Mets at Dx, continued

5. If the only indication of extension in the record is the physician's statement of an M category from the TNM staging system or a stage from a site-specific staging system, such as Dukes' D, record the numerically lowest equivalent extension code for that M category. In most cases, this will be 40, Distant metastasis, NOS.
6. If the information in the medical record is ambiguous or incomplete regarding the extent to which the tumor has spread, the extent of disease may be inferred from the M category stated by the physician.
7. Some site or histology schemas include a designation of M1, NOS. The NOS is added when there is further breakdown of the category into subsets (such as M1a, M1b, M1c), but the correct subset cannot be determined. The NOS designation, which can appear in both the descriptions of codes and the mapping, is not official AJCC descriptive terminology. The NOS should be disregarded in reports and analyses when it is not a useful distinction. The data collector should only code to a category such as "Stated as M1 NOS" when the appropriate subset (such as M1a or M1b) cannot be determined.
8. It is strongly recommended that the choice of distant lymph nodes and/or distant metastasis codes be documented in a related text field on the abstract.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS METS EVAL

Item Length: 2

NAACCR Item #2860

Description

Records how the code for the item “CS Mets at Dx” was determined based on the diagnostic methods employed.

Code	Description	Staging Basis
0	No pathologic examination of metastatic tissue performed. Evaluation of distant metastasis based on physical examination, imaging examination, and/or other non-invasive clinical evidence. No autopsy evidence used.	c
1	No pathologic examination of metastatic tissue performed. Evaluation of distant metastasis based on endoscopic examination or other invasive technique, including surgical observation without biopsy. No autopsy evidence used. <i>Does not meet criteria for AJCC pathologic staging of distant metastasis.</i>	c
2	No pathologic examination of metastatic tissue done prior to death, but evidence derived from autopsy (tumor was suspected or diagnosed prior to autopsy).	p
3	Pathologic examination of metastatic tissue performed WITHOUT pre-surgical systemic treatment or radiation OR pathologic examination of metastatic tissue performed, unknown if pre-surgical systemic treatment or radiation performed <i>Meets criteria for AJCC pathologic staging of distant metastasis.</i>	p
5	Pathologic examination of metastatic tissue performed WITH pre-surgical systemic treatment or radiation, and metastasis based on clinical evidence.	c
6	Pathologic examination of metastatic tissue performed WITH pre-surgical systemic treatment or radiation, BUT metastasis based on pathologic evidence.	y
8	Evidence from autopsy AND tumor was unsuspected or undiagnosed prior to autopsy.	a
9	Not assessed; cannot be assessed Unknown if assessed Not documented in patient record <i>For sites with no TNM staging:</i> Not applicable	c

Instructions for Coding

1. One of the uses of the CS Mets Eval field is to assign a “c” or “p” to the M category derived from the CS Mets at DX field. Since both clinical and pathologic evidence might be available for assessing distant metastasis, the coding of the Eval field can be confusing. The goal is to assign the Eval code that indicates the best evidence used to determine the M category. Coding of the Eval field thus requires that the abstractor take note of the M category that will be derived from the code in the CS Mets at DX field and then use the following guidelines to determine the best Eval code to assign.
 - a. If M0 will be derived (i.e., no distant metastasis are coded), then choose an Eval code that will derive a “c” staging basis. There is no category of pM0, because it is impossible to disprove all possible sites of metastasis pathologically.
Example Cecum carcinoma with negative chest X-ray and negative liver biopsy. CS Mets at DX is coded 00 (None), which maps to M0. CS Mets Eval is coded 1 (Evaluation of distant metastasis based on endoscopic examination or other invasive technique,

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Mets Eval, continued

- including surgical observation without biopsy), which maps to the “c” staging basis.
- b. If MX will be derived (i.e., CS Mets at DX is coded 99), then choose an Eval code that will derive a “c” staging basis. The appropriate code might be 9 (Unknown) or might be another code if workup was done but the results were not definitively positive or negative.

Example Cecum carcinoma abstracted from a pathology report of biopsy only, no clinical data or surgical observations available, CS Mets at DX coded 99 (Unknown) which will map to MX. CS Mets Eval is coded 9 (Unknown), which maps to the “c” staging basis.

Example Lung cancer diagnosed by imaging. Patient has behavior changes, and brain imaging cannot rule out metastases. Patient is not a surgical candidate. CS Mets at DX is coded 99 (Unknown) which maps to MX. CS Mets Eval is coded 0 (imaging), which maps to the “c” staging basis.

- c. If M1 will be derived (i.e., there is disease present that is coded in the CS Mets at DX field) and there are no subcategories of M1, such as M1a and M1b, then determine if there was any pathological evidence for the M1 category. If so, select an Eval code that will derive a “p” staging basis. If there was only clinical evidence of the M1 disease, select an Eval code that will derive a “c” staging basis.

Example Cecum carcinoma with negative chest X-ray and positive liver biopsy. CS Mets at DX is coded 40 (Distant metastasis except distant nodes), which maps to M1, and there are no subcategories of M for the colon schema. CS Mets Eval is coded 3 (Pathologic examination of metastatic tissue performed WITHOUT pre-surgical systemic treatment or radiation OR pathologic examination of metastatic tissue performed, unknown if pre-surgical systemic treatment or radiation performed), which maps to the “p” staging basis.

Example Cecum carcinoma with positive chest X-ray and negative liver biopsy. CS Mets at DX is coded 40 (Distant metastasis except distant nodes), which maps to M1, and there are no subcategories of M for the colon schema. CS Mets Eval is coded 0 (No pathologic examination of metastatic tissue performed. Evaluation based on physical examination, imaging examination, and/or other non-invasive clinical evidence. No autopsy evidence used), which maps to the “c” staging basis.

- d. If a specific subcategory of M1 will be derived (such as M1a, etc.), then determine if there was any pathological evidence for the specific subcategory. If so, select an Eval code that will derive a “p” staging basis. If there was only clinical evidence of the subcategory disease, select an Eval code that will derive a “c” staging basis. In the latter case there may have been pathological evidence of a lower M subcategory, but this is not considered in assigning the Eval code.

Example Prostate carcinoma with the following:

Involvement	CS Mets at DX Code	TNM
Positive biopsy of aortic lymph node (distant node)	Code 12	pM1a
Positive bone imaging	Code 30	cM1b
Positive brain imaging	Code 40	cM1c
All of the above	Codes 12 + 30 + 40 = Code 55	cM1c

To code CS Mets at DX, follow the general rule to code the highest applicable code, even though there is pathological evidence of metastases. CS Mets at DX is coded 55, which combines the codes for the lymph node, bone, and brain involvement. Code 55

Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items

CS Mets Eval, continued

maps to M1c. There is no pathological evidence for the subcategory of M1c (the only pathological evidence is for subcategory M1a). CS Mets Eval is coded 0 (imaging), which maps to the “c” staging basis. The positive lymph node would map to M1a, a lower M subcategory, so do not base the Eval code on that.

Example Prostate carcinoma with positive biopsy of aortic lymph node (distant node), negative bone scan, and negative brain scan. CS Mets at DX is coded 12 (distant lymph node), which maps to M1a. CS Mets Eval is coded 3 (Pathologic examination of metastatic tissue performed WITHOUT pre-surgical systemic treatment or radiation OR pathologic examination of metastatic tissue performed, unknown if pre-surgical systemic treatment or radiation performed), which maps to the “p” staging basis.

2. For sites and histologies for which no TNM schema has been defined, such as brain or Kaposi sarcoma, this field is always coded 9, Not Applicable. (See Table 6 in the General Instructions.) For any sites and histologies not listed in Table 6, code to the value that best reflects the diagnostic methods used, whether or not a stage is actually calculated for an individual case. In other words, do not use code 9 when a case has a histology that is excluded from staging but the site does have a TNM schema defined, for example, a sarcoma of the breast. In those cases, use code 9 only when the nature of the diagnostic methods is actually unknown.
3. If the patient receives preoperative (neoadjuvant) systemic therapy (chemotherapy, hormone therapy, immunotherapy) or radiation therapy, the clinical status of metastases at diagnosis takes precedence (code 5), unless the pathologic evidence is more extensive (code 6).
4. Code 0 includes imaging studies such as standard radiography, special radiographic projections, tomography, computerized tomography (CT), ultrasonography (US), lymphography, angiography, scintigraphy (nuclear scans), magnetic resonance imaging (MRI), positron emission tomography (PET), spiral scanning (CT or MRI) and other non-invasive methods of examining tissues.
5. Code 1 includes endoscopy and observations at surgery, such as abdominal exploration at the time of a colon resection, where distant metastasis is not biopsied.
6. AJCC does not recognize a pathologic M0 category since it is not possible to rule out all possible metastatic sites. Therefore, if the patient has a biopsy or removal of a distant site and the pathology report is negative, generally use Eval code 1, because this does not meet the criteria for pathologic staging.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

Page left blank.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS SITE-SPECIFIC FACTOR 1

Item Length: 3

NAACCR Item #2880

Description

Identifies additional information needed to generate stage, or prognostic factors that have an effect on stage or survival.

Code	Description
000	None
	SITE/HISTOLOGY-SPECIFIC CODES
999	Unknown; [site-specific title] cannot be assessed; Not documented in patient record

For schemas that do not use this site-specific factor:

Code	Description
888	Not applicable for this site

Instructions for Coding

1. If there is no site/histology-specific factor for a schema, code 888.
2. The following primary sites/histologies use Site Specific Factor 1 to code information. See the site-specific schemas for acceptable codes and their definitions.

<u>Site/Histology</u>	<u>Factor</u>
Head and neck*	Size of Lymph Nodes
Colon	Carcinoembryonic Antigen (CEA)
Rectosigmoid, rectum	Carcinoembryonic Antigen (CEA)
Liver	Alpha Fetoprotein (AFP)
Pleura	Pleural Effusion
Malignant Melanoma of Skin, Vulva, Penis, Scrotum	Measured Thickness (Depth), Breslow's Measurement
Mycosis Fungoides	Peripheral Blood Involvement
Breast	Estrogen Receptor Assay (ERA)
Ovary	Carbohydrate Antigen 125 (CA-125)
Placenta	Prognostic Scoring Index
Prostate	Prostate Specific Antigen Laboratory Value (PSA PSA Lab Value)
Testis	Alpha Fetoprotein (AFP)
Thyroid	Single vs. Multiple Nodules

* Head and neck includes the following schemas: upper lip; lower lip; other lip; base of tongue; anterior tongue; upper gum; lower gum and retromolar trigone; other gum; floor of mouth; hard palate; soft palate/uvula; other mouth; buccal mucosa; parotid gland; submandibular gland; other salivary glands; oropharynx; anterior surface of epiglottis; nasopharynx; pyriform sinus/hypopharynx; other pharynx; nasal cavity; middle ear; maxillary sinus; ethmoid sinus; other sinus; glottic larynx; supraglottic larynx; subglottic larynx; other larynx

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Site-Specific Factor 1, continued

<u>Site/Histology</u>	<u>Factor</u>
Melanoma of Conjunctiva	Measured Thickness (Depth), Breslow's Measurement
Melanoma of Choroid	Measured Thickness (Depth), Breslow's Measurement
Melanoma of Iris and Ciliary Body	Measured Thickness (Depth), Breslow's Measurement
Retinoblastoma	Extension Evaluated at Enucleation
Brain	WHO Grade
Other CNS	WHO Grade
Other Endocrine	WHO Grade
Kaposi Sarcoma	Associated with HIV/AIDS
Lymphoma	Associated with HIV/AIDS

3. Code 000 Not done is used when there is a statement in the record that a test was not performed.
 - a. If there is no report of a lab test in the patient record, code as 999 Unknown; Not documented in patient record.
 - b. For Kaposi sarcoma, if AIDS status is not documented, code as 999 Unknown rather than 002, Not Present.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS SITE-SPECIFIC FACTOR 2

Item Length: 3

NAACCR Item #2890

Description

Identifies additional information needed to generate stage, or prognostic factors that have an effect on stage or survival.

Code	Description
000	None
	SITE/HISTOLOGY-SPECIFIC CODES
999	Unknown; [site-specific title] cannot be assessed; Not documented in patient record

For schemas that do not use this site-specific factor:

Code	Description
888	Not applicable for this site

Instructions for Coding

1. If there is no site/histology-specific factor for a schema, code 888.
2. The following primary sites use Site Specific Factor 2 to code information. See the site-specific schemas for acceptable codes and their definitions.

Site/Histology

Head and neck*

Liver

Malignant Melanoma of Skin,

Vulva, Penis, Scrotum

Breast

Prostate

Testis

Hodgkin and non-Hodgkin Lymphoma

Factor

Extracapsular Extension, Lymph Nodes for
Head and Neck

Fibrosis Score

Ulceration

Progesterone Receptor Assay (PRA)

Prostate Specific Antigen (PSA)

Human Chorionic Gonadotropin (HCG)

Symptoms at Diagnosis

* Head and neck includes the following schemas: upper lip; lower lip; other lip; base of tongue; anterior tongue; upper gum; lower gum and retromolar trigone; other gum; floor of mouth; hard palate; soft palate/uvula; other mouth; buccal mucosa; parotid gland; submandibular gland; other salivary glands; oropharynx; anterior surface of epiglottis; nasopharynx; pyriform sinus/hypopharynx; other pharynx; nasal cavity; middle ear; maxillary sinus; ethmoid sinus; other sinus; glottic larynx; supraglottic larynx; subglottic larynx; other larynx

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Site-Specific Factor 2, continued

3. Code 000 Not done is used when there is a statement in the record that a test was not performed.
 - a. If there is no report of a lab test in the health record, code as 999 Unknown; Not documented in patient record.
 - b. For malignant melanoma of skin, if ulceration is not mentioned in the pathology report, code as 000 No ulceration present.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS SITE-SPECIFIC FACTOR 3

Item Length: 3

NAACCR Item #2900

Description

Identifies additional information needed to generate stage, or prognostic factors that have an effect on stage or survival.

Code	Description
000	None
	SITE/HISTOLOGY-SPECIFIC CODES
999	Unknown; [site-specific title] cannot be assessed; Not documented in patient record

For schemas that do not use this site-specific factor:

Code	Description
888	Not applicable for this site

Instructions for Coding

- If there is no site/histology-specific factor for a schema, code 888.
- The following primary sites use Site Specific Factor 3 to code information. See the site-specific schemas for acceptable codes and their definitions.

Site/Histology

Head and Neck*
Malignant Melanoma of Skin,
 Vulva, Penis, Scrotum
Breast
Prostate
Testis
Lymphoma

Factor

Levels I-III, Lymph Nodes of Head and Neck

Clinical Status of Lymph Node Mets
Number of Positive Ipsilateral Axillary Lymph Nodes
CS Extension - Pathologic Extension
LDH (Lactate Dehydrogenase)
International Prognostic Index (IPI) Score

* Head and neck includes the following schemas: upper lip; lower lip; other lip; base of tongue; anterior tongue; upper gum; lower gum and retromolar trigone; other gum; floor of mouth; hard palate; soft palate/uvula; other mouth; buccal mucosa; parotid gland; submandibular gland; other salivary glands; oropharynx; anterior surface of epiglottis; nasopharynx; pyriform sinus/hypopharynx; other pharynx; nasal cavity; middle ear; maxillary sinus; ethmoid sinus; other sinus; glottic larynx; supraglottic larynx; subglottic larynx; other larynx

- Code 000 Not done is used when there is a statement in the record that a test was not performed.
 - If there is no report of a lab test in the health record, code as 999 Unknown; Not documented in patient record.
 - For the lymphomas, if the IPI score is not stated in the record, code as 999 Unknown; Not documented in patient record. It is not necessary to calculate the IPI score from other information in the record.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Site-Specific Factor 3, continued

For head and neck sites only:

4. Use code 9 only when it is unknown if lymph nodes are involved. Within the Site-Specific Factors, do not code 9 in some positions and 0 or 1 in other positions. If specific information is available about the positive or negative status of some but not all nodes in any one level or group, assume that the rest of the nodes in the same Site-Specific Factor are negative and code accordingly.
5. When the only information available is “Regional nodes, NOS” or “Cervical nodes, NOS” or “Internal jugular lymph nodes, NOS” or “Lymph nodes, NOS,” code 0 in all digits of Site-Specific Factors 3-6.
6. See “Coding Regional Lymph Nodes for Head and Neck Sites” under CS Lymph Nodes for further information about the regional nodes of the head and neck, including definitions of the levels.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS SITE-SPECIFIC FACTOR 4

Item Length: 3

NAACCR Item #2910

Description

Identifies additional information needed to generate stage, or prognostic factors that have an effect on stage or survival.

Code	Description
000	None
	SITE/HISTOLOGY-SPECIFIC CODES
999	Unknown; [site-specific title] cannot be assessed; Not documented in patient record

For schemas that do not use this site-specific factor:

Code	Description
888	Not applicable for this site

Instructions for Coding

1. If there is no site/histology-specific factor for a schema, code 888.
2. The following primary sites use Site Specific Factor 4 to code information. See the site-specific schemas for acceptable codes and their definitions.

Site/Histology

Head and Neck*
Malignant Melanoma of Skin,
 Vulva, Penis, Scrotum
Breast
Prostate

Factor

Levels IV-V, Lymph Nodes of Head and Neck

Lactate Dehydrogenase (LDH)
Immunohistochemistry (IHC) of Regional Lymph Nodes
Prostate Apex Involvement (effective as of version 1.02.00)
[Prostatic Acid Phosphatase (PAP)–OBSOLETE as of version 1.02.00]

Radical Orchiectomy Performed

Testis

* Head and neck includes the following schemas: upper lip; lower lip; other lip; base of tongue; anterior tongue; upper gum; lower gum and retromolar trigone; other gum; floor of mouth; hard palate; soft palate/uvula; other mouth; buccal mucosa; parotid gland; submandibular gland; other salivary glands; oropharynx; anterior surface of epiglottis; nasopharynx; pyriform sinus/hypopharynx; other pharynx; nasal cavity; middle ear; maxillary sinus; ethmoid sinus; other sinus; glottic larynx; supraglottic larynx; subglottic larynx; other larynx

3. Code 000 Not done is used when there is a statement in the record that a test was not performed.
 - a. If there is no report of a lab test in the health record, code as 999 Unknown; Not documented in patient record.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Site-Specific Factor 4, continued

For head and neck sites only:

4. Use code 9 only when it is unknown if lymph nodes are involved. Within the Site-Specific Factors, do not code 9 in some positions and 0 or 1 in other positions. If specific information is available about the positive or negative status of some but not all nodes in any one level or group, assume that the rest of the nodes in the same Site-Specific Factor are negative and code accordingly.
5. When the only information available is “Regional nodes, NOS” or “Cervical nodes, NOS” or “Internal jugular lymph nodes, NOS” or “Lymph nodes, NOS,” code 0 in all digits of Site-Specific Factors 3-6.
6. See “Coding Regional Lymph Nodes for Head and Neck Sites” under CS Lymph Nodes for further information about the regional nodes of the head and neck, including definitions of the levels.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS SITE-SPECIFIC FACTOR 5

Item Length: 3

NAACCR Item #2920

Description

Identifies additional information needed to generate stage, or prognostic factors that have an effect on stage or survival.

Code	Description
000	None
	SITE/HISTOLOGY-SPECIFIC CODES
999	Unknown; [site-specific title] cannot be assessed; Not documented in patient record

For schemas that do not use this site-specific factor:

Code	Description
888	Not applicable for this site

Instructions for Coding

1. If there is no site/histology-specific factor for a schema, code 888.
2. The following primary sites use Site Specific Factor 5 to code information. See the site-specific schemas for acceptable codes and their definitions.

Site/Histology

Head and Neck*
Breast
Prostate
Testis

Factor

Levels VI-VIII, Lymph Nodes of Head and Neck
Molecular Studies of Regional Lymph Nodes
Gleason's Primary and Secondary Patterns
Size of Metastasis in Lymph Nodes

* Head and neck includes the following schemas: upper lip; lower lip; other lip; base of tongue; anterior tongue; upper gum; lower gum and retromolar trigone; other gum; floor of mouth; hard palate; soft palate/uvula; other mouth; buccal mucosa; parotid gland; submandibular gland; other salivary glands; oropharynx; anterior surface of epiglottis; nasopharynx; pyriform sinus/hypopharynx; other pharynx; nasal cavity; middle ear; maxillary sinus; ethmoid sinus; other sinus; glottic larynx; supraglottic larynx; subglottic larynx; other larynx

3. Code 000 Not done is used when there is a statement in the record that a test was not performed.
 - a. If there is no report of a lab test in the health record, code as 999 Unknown; Not documented in patient record.

For head and neck sites only:

4. Use code 9 only when it is unknown if lymph nodes are involved. Within the Site-Specific Factors, do not code 9 in some positions and 0 or 1 in other positions. If specific information is available about the positive or negative status of some but not all nodes in any one level or group, assume that the rest of the nodes in the same Site-Specific Factor are negative and code accordingly.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Site-Specific Factor 5, continued

5. When the only information available is “Regional nodes, NOS” or “Cervical nodes, NOS” or “Internal jugular lymph nodes, NOS” or “Lymph nodes, NOS,” code 0 in all digits of Site-Specific Factors 3-6.

6. See “Coding Regional Lymph Nodes for Head and Neck Sites” under CS Lymph Nodes for further information about the regional nodes of the head and neck, including definitions of the levels.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS SITE-SPECIFIC FACTOR 6

Item Length: 3

NAACCR Item #2930

Description

Identifies additional information needed to generate stage, or prognostic factors that have an effect on stage or survival.

Code	Description
000	None
	SITE/HISTOLOGY-SPECIFIC CODES
999	Unknown; [site-specific title] cannot be assessed; Not documented in patient record

For schemas that do not use this site-specific factor:

Code	Description
888	Not applicable for this site

Instructions for Coding

1. If there is no site/histology-specific factor for a schema, code 888.
2. The following primary sites use Site Specific Factor 6 to code information. See the site-specific schemas for acceptable codes and their definitions.

Site/Histology

Head and Neck*

Breast

Prostate

Factor

Parapharyngeal, Parotid, Preauricular, and Sub-Occipital Lymph Nodes, Lymph Nodes for Head and Neck

Size of Tumor--Invasive Component

Gleason's Score

* Head and neck includes the following schemas: upper lip; lower lip; other lip; base of tongue; anterior tongue; upper gum; lower gum and retromolar trigone; other gum; floor of mouth; hard palate; soft palate/uvula; other mouth; buccal mucosa; parotid gland; submandibular gland; other salivary glands; oropharynx; anterior surface of epiglottis; nasopharynx; pyriform sinus/hypopharynx; other pharynx; nasal cavity; middle ear; maxillary sinus; ethmoid sinus; other sinus; glottic larynx; supraglottic larynx; subglottic larynx; other larynx

3. Code 000 Not done is used when there is a statement in the record that a test was not performed.
 - a. If there is no report of a lab test in the health record, code as 999 Unknown; Not documented in patient record.

**Collaborative Staging Manual and Coding Instructions Part I
Instructions for Data Items**

CS Site-Specific Factor 6, continued

For head and neck sites only:

4. Use code 9 only when it is unknown if lymph nodes are involved. Within the Site-Specific Factors, do not code 9 in some positions and 0 or 1 in other positions. If specific information is available about the positive or negative status of some but not all nodes in any one level or group, assume that the rest of the nodes in the same Site-Specific Factor are negative and code accordingly.
5. When the only information available is “Regional nodes, NOS” or “Cervical nodes, NOS” or “Internal jugular lymph nodes, NOS” or “Lymph nodes, NOS,” code 0 in all digits of Site-Specific Factors 3-6.
6. See “Coding Regional Lymph Nodes for Head and Neck Sites” under CS Lymph Nodes for further information about the regional nodes of the head and neck, including definitions of the levels.

Collaborative Staging Manual and Coding Instructions Part I Additional Tables and Appendices

ADDITIONAL COLLABORATIVE STAGING TABLES

In addition to the tables of codes for each of the Collaborative Staging System data items, it was necessary to develop reference tables that the computer algorithm uses to assure that the output data items (T, N, M, Stage Group, SS77 and SS2000) are accurately derived. These tables are not printed in this manual, usually because of their length. Any reference tables that have been developed for individual schema are listed at the top of the schema in Part II of this manual but are not printed in the manual. They are available for reference on the Collaborative Staging website, www.cancerstaging.org.

AJCC STAGE TABLE

The allowable storage codes for derived T, N, M and Stage Group are shown in Appendix 2 with their output character strings. The AJCC stage tables are site-specific and are not included in this manual due to their length. The data collector or researcher can access the AJCC stage table associated with each schema under the appropriate site-specific section of the Collaborative Staging website, www.cancerstaging.org.

SUMMARY STAGE TABLE

The summary stage conversion table is shown in Appendix 3. This table evaluates the CS Extension, CS Lymph Nodes, and CS Mets at Dx fields to determine the final Summary Stage 77 and Summary Stage 2000 output (Appendix 2e). The Summary Stage Table applies to all schemas and lists all possible combinations, including Not Applicable, Unstagable and Error situations. The algorithm takes the highest (most extensive value) from any of the three input fields as the output value. For example, if the Extension code maps to regional direct extension, the Lymph Nodes code maps to regional lymph nodes, and no distant metastases are coded in the Mets at Dx field, the output value will be RE+RN, regional extension and nodes.

HISTOLOGY EXCLUSION TABLES

It has been previously noted that not all cases will have T, N, M, and Stage Group categories derived by the computer algorithm. This is because certain histologies are excluded from some chapters of the *AJCC Cancer Staging Manual, sixth edition*. In order that the Collaborative Staging System could accurately derive the components of the TNM system only for the histologies allowed in the AJCC manual, tables of allowable and excluded ICD-O-3 histology have been developed with the cooperation of the AJCC. These lists are shown in Appendix 5 of this manual. The nine major categories of cancers are shown with their associated three-digit morphology code ranges.

For example, in TNM staging, carcinoids are specifically excluded from the colon cancer chapter. If a malignant carcinoid case is abstracted, all of the 10 data items for colon should be recorded (9 basic data items plus one site-specific factor for colon). The computer algorithm will look at the recorded ICD-O-3 morphology coded and match it to the exclusions table for colon. Because carcinoid (M-8240/3) is on the exclusions list, the algorithm will not generate a T, N, M or Stage Group, but will generate both Summary Stage 1977 and 2000.

SITE-SPECIFIC EXTRA TABLES

In the introduction to this manual it was noted that some schemas require additional reference tables in order for the computer algorithm to determine the final derived T, N, M, or Stage Group output. The need for these extra tables arises when additional information is needed to differentiate, for example, a T1a from a T1b, or when the tumor size is a significant factor in determining the T category.

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

For example, it is necessary to combine information from both the tumor size table and the extension table in order to derive the T category for breast cancer. If the tumor extension is purely in situ, the derived T is Tis; if the tumor extension involves the skin or chest wall, the derived T is one of the T4 subcategories. But if the tumor extension is in the range of 10-30, it is necessary to know the exact size of the tumor. The computer algorithm looks at the “Extension Size Table” for breast to determine the correct output. In the table below, if the Extension code is 10 and the tumor size is coded 018 (1.8 cm), the computer algorithm will read the sixth line of the table and output a T1c. If the extension code is 20 and the tumor size is coded 055, the computer algorithm will read the eighth line of the table and output a T3.

Figure 3. Example of “Extension Size Table” for Breast Schema

Note: For Extension codes 10, 20, and 30 ONLY, the T category is assigned based on value of CS Tumor Size, as follows:

From Tumor Size	To Tumor Size	T Code	Comment
000	000	ERROR	Tumor size 000 should only be used with Extension 95.
001	001	T1mic	
002	005	T1a	
006	010	T1b	
011	020	T1c	
021	050	T2	
051	989	T3	
990	990	T1mic	
991	991	T1b	
992	992	T1c	
993	995	T2	
996	996	T1NOS	Per downstaging rule.
997	997	ERROR	Tumor size 997 should only be used with Extension code 05 or 07.
998	998	T3	
999	999	TX	

As another example, the patient’s age and histology must be known in order to stage a thyroid cancer. Several additional tables on the thyroid schema are used by the computer algorithm to determine the TNM Stage Group when the patient is under or over age 45 and the histology is papillary/follicular, medullary or anaplastic.

Appendices

1. Determining Descriptive Tumor Size
2. Output Values, Storage Codes, and Display String Descriptions for T, N, M, Stage Group and Summary Stage
3. Summary Stage Conversion Algorithm for All Schemas
4. Site Specific Factors
5. Histology Exclusion Groups

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

**Appendix 1.
Determining Descriptive Tumor Size
Millimeter Equivalents for Descriptive Terms**

<u>Fruits</u>	<u>mm</u>	<u>Miscellaneous Food</u>	<u>mm</u>
Apple	070	Doughnut	090
Apricot	040	Egg	050
Cherry	020	Bantam	040
Date	040	Goose	070
Fig (dried)	040	Hen	030
Grape	020	Pigeon	030
Grapefruit	100	Robin	020
Kumquat	050	Lentil	991
Lemon	080	Millet	991
Olive	020		
Orange	090	<u>Money</u>	
Peach	060	Dime	010
Pear	090	Dollar, half	030
Plum	030	Dollar, silver	040
Tangerine	060	Nickel	020
		Penny	010
<u>Nuts</u>		Quarter	020
Almond	030	Penny	010
Chestnut	040		
Chestnut, horse	040	<u>Other</u>	
Hazel	020	Ball, golf	040
Hickory	030	Ball, ping-pong	030
Peanut	010	Ball, tennis	060
Pecan	030	Baseball	070
Walnut	030	Eraser on pencil	991
		Fist	090
<u>Vegetables</u>		Marble	010
Bean	010	Matchhead	991
Bean, lima	020		
Pea	991	Microscopic focus	990
Pea, split	991		

SIZES IN CENTIMETERS, MILLIMETERS, INCHES

10 millimeters (mm) = 1 centimeter (cm)

1 millimeter (mm) = 1/10 centimeter (cm)

2.5 centimeters (cm) = 1 inch (in)

1 centimeter (cm) = .394 inch (in)

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

APPENDIX 2. ALLOWABLE VALUES

Appendix 2a. T Allowable Codes

This table shows the allowable values for the generated Collaborative Stage data items. The Storage Code is the value to be stored in the field of a NAACCR record. The Storage Codes are designed for analysis. The Display String is the corresponding label that should be displayed on the screen or in a report. The meaning of these strings will be clear to the registrar or physician user.

Storage Code	Display String*	Display String Description
99	TX	TX
00	T0	T0
01	Ta	Ta
05	Tis	Tis
06	Tispu	Tispu (Urethra only)
07	Tispd	Tispd (Urethra only)
10	T1	T1
11	T1mic	T1mic
12	T1a	T1a
13	T1a1	T1a1
14	T1a2	T1a2
15	T1b	T1b
16	T1b1	T1b1
17	T1b2	T1b2
18	T1c	T1c
19	T1NOS	T1 NOS
20	T2	T2
29	T2NOS	T2 NOS
21	T2a	T2a
22	T2b	T2b
23	T2c	T2c
30	T3	T3
39	T3NOS	T3 NOS
31	T3a	T3a
32	T3b	T3b
33	T3c	T3c
40	T4	T4
49	T4NOS	T4 NOS
41	T4a	T4a
42	T4b	T4b
43	T4c	T4c
44	T4d	T4d
88	NA	Not applicable

* T_ with no subscript indicates that there is only one choice for that category
T_ NOS indicates that there are additional choices for the category but a more specific code cannot be determined.

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Appendix 2b. N Allowable Codes

This table shows the allowable values for the generated Collaborative Stage data items. The Storage Code is the value to be stored in the field of a NAACCR record. The Storage Codes are designed for analysis. The Display String is the corresponding label that should be displayed on the screen or in a report. The meaning of these strings will be clear to the registrar or physician user.

Storage Code	Display String*	Display String Description
99	NX	NX
00	N0	N0
09	N0NOS	N0 NOS
01	N0(i-)	N0(i-)
02	N0(i+)	N0(i+)
03	N0(mol-)	N0(mol-)
04	N0(mol+)	N0(mol+)
10	N1	N1
19	N1NOS	N1 NOS
11	N1a	N1a
12	N1b	N1b
13	N1c	N1c
18	N1mi	N1mi
20	N2	N2
29	N2NOS	N2 NOS
21	N2a	N2a
22	N2b	N2b
23	N2c	N2c
30	N3	N3
39	N3NOS	N3 NOS
31	N3a	N3a
32	N3b	N3b
33	N3c	N3c
88	NA	Not applicable

* N_ with no subscript indicates that there is only one choice for that category

N_ NOS indicates that there are additional choices for the category but a more specific code cannot be determined.

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Appendix 2c. M Allowable Codes

This table shows the allowable values for the generated Collaborative Stage data items. The Storage Code is the value to be stored in the field of a NAACCR record. The Storage Codes are designed for analysis. The Display String is the corresponding label that should be displayed on the screen or in a report. The meaning of these strings will be clear to the registrar or physician user.

Storage Code	Display String*	Display String Description
99	MX	MX
00	M0	M0
10	M1	M1
11	M1a	M1a
12	M1b	M1b
13	M1c	M1c
19	M1NOS	M1 NOS
88	NA	Not applicable

* M_ with no subscript indicates that there is only one choice for that category

M_ NOS indicates that there are additional choices for the category but a more specific code cannot be determined.

Appendix 2d. Stage Allowable Codes

This table shows the allowable values for the generated Collaborative Stage data items. The Storage Code is the value to be stored in the field of a NAACCR record. The Storage Codes are designed for analysis. The Display String is the corresponding label that should be displayed on the screen or in a report. The meaning of these strings will be clear to the registrar or physician user.

Storage Code	Display String*	Display String Description
00	0	Stage 0
01	0a	Stage 0a
02	0is	Stage 0is
10	I	Stage I
11	INOS	Stage I NOS
12	IA	Stage IA
13	IA1	Stage IA1
14	IA2	Stage IA2
15	IB	Stage IB
16	IB1	Stage IB1
17	IB2	Stage IB2
18	IC	Stage IC
19	IS	Stage IS
23	ISA	Stage ISA (lymphoma only)
24	ISB	Stage ISB (lymphoma only)
20	IEA	Stage IEA (lymphoma only)
21	IEB	Stage IEB (lymphoma only)
22	IE	Stage IE (lymphoma only)
30	II	Stage II

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

**Appendix 2d
continued
Storage Code**

Storage Code	Display String	Display String Description
31	IINOS	Stage II NOS
32	IIA	Stage IIA
33	IIB	Stage IIB
34	IIC	Stage IIC
35	IIEA	Stage IIEA (lymphoma only)
36	IIEB	Stage IIEB (lymphoma only)
37	IIE	Stage IIE (lymphoma only)
38	IISA	Stage IISA (lymphoma only)
39	IISB	Stage IISB (lymphoma only)
40	IIS	Stage IIS (lymphoma only)
41	IIESA	Stage IIESA (lymphoma only)
42	IIESB	Stage IIESB (lymphoma only)
43	IIES	Stage IIES (lymphoma only)
50	III	Stage III
51	IIINOS	Stage III NOS
52	IIIA	Stage IIIA
53	IIIB	Stage IIIB
54	IIIC	Stage IIIC
55	IIIEA	Stage IIIEA (lymphoma only)
56	IIIEB	Stage IIIEB (lymphoma only)
57	IIIE	Stage IIIE (lymphoma only)
58	IIISA	Stage IIISA (lymphoma only)
59	IIISB	Stage IIISB (lymphoma only)
60	IIIS	Stage IIIS (lymphoma only)
61	IIIESA	Stage IIIESA (lymphoma only)
62	IIIESB	Stage IIIESB (lymphoma only)
63	IIIES	Stage IIIES (lymphoma only)
70	IV	Stage IV
71	IVNOS	Stage IV NOS
72	IVA	Stage IVA
73	IVB	Stage IVB
74	IVC	Stage IVC
88	NA	Not applicable
90	OCCULT	Stage Occult
99	UNK	Stage Unknown

* A stage group with no subscript indicates that there is only one choice for that category
A stage group NOS indicates that there are additional choices for the category but a more specific code cannot be determined.

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Appendix 2e. Summary Stage Allowable Codes

This table shows the allowable values for the generated Collaborative Stage data items. The Storage Code is the value to be stored in the field of a NAACCR record. The Storage Codes are designed for analysis. The Display String is the corresponding label that should be displayed on the screen or in a report. The meaning of these strings will be clear to the registrar or physician user.

Storage Code	Display String	Display String Description
	ERROR	Processing error (no storage code needed)
	NONE	None (internal use only, no storage code needed)
0	IS	In situ
1	L	Localized
2	RE	Regional, direct extension
3	RN	Regional, lymph nodes only
4	RE+RN	Regional, extension and nodes
5	RNOS	Regional, NOS
7	D	Distant
8	NA	Not applicable
9	U	Unknown/Unstaged

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Appendix 3. Summary Stage Conversion Algorithm for All Schemas

* In situ implies no involvement outside the primary site.

Extension SS77 or SS2000 result	LN SS77 or SS2000 result	Mets SS77 or SS2000 result	Final SS77 or SS2000 result	Extension SS77 or SS2000 result	LN SS77 or SS2000 result	Mets SS77 or SS2000 result	Final SS77 or SS2000 result	Extension SS77 or SS2000 result	LN SS77 or SS2000 result	Mets SS77 or SS2000 result	Final SS77 or SS2000 result
IS*	D	RE+RN	ERROR	L	RN	RE	RE+RN				
IS*	D	D	ERROR	L	RN	RN	RN				
IS*	NONE	NONE	IS	IS*	D	NA	ERROR	L	RN	RE+RN	RE+RN
IS*	NONE	L	ERROR	IS*	D	U	ERROR	L	RN	D	D
IS*	NONE	RE	ERROR	IS	NA	NONE	IS	L	RN	NA	RN
IS*	NONE	RN	ERROR	IS*	NA	L	ERROR	L	RN	U	RN
IS*	NONE	RE+RN	ERROR	IS*	NA	RE	ERROR	L	RE+RN	NONE	RE+RN
IS*	NONE	D	ERROR	IS*	NA	RN	ERROR	L	RE+RN	L	RE+RN
IS	NONE	NA	IS	IS*	NA	RE+RN	ERROR	L	RE+RN	RE	RE+RN
IS	NONE	U	IS	IS*	NA	D	ERROR	L	RE+RN	RN	RE+RN
IS*	RE	NONE	ERROR	IS	NA	NA	IS	L	RE+RN	RE+RN	RE+RN
IS*	RE	L	ERROR	IS	NA	U	IS	L	RE+RN	D	D
IS*	RE	RE	ERROR	IS*	U	NONE	IS	L	RE+RN	NA	RE+RN
IS*	RE	RN	ERROR	IS*	U	L	ERROR	L	RE+RN	U	RE+RN
IS*	RE	RE+RN	ERROR	IS*	U	RE	ERROR	L	D	NONE	D
IS*	RE	D	ERROR	IS*	U	RN	ERROR	L	D	L	D
IS*	RE	NA	ERROR	IS*	U	RE+RN	ERROR	L	D	RE	D
IS*	RE	U	ERROR	IS*	U	D	ERROR	L	D	RN	D
IS*	RN	NONE	ERROR	IS	U	NA	IS	L	D	RE+RN	D
IS*	RN	L	ERROR	IS	U	U	IS	L	D	D	D
IS*	RN	RE	ERROR	L	NONE	NONE	L	L	D	NA	D
IS*	RN	RN	ERROR	L	NONE	L	L	L	D	U	D
IS*	RN	RE+RN	ERROR	L	NONE	RE	RE	L	NA	NONE	L
IS*	RN	D	ERROR	L	NONE	RN	RN	L	NA	L	L
IS*	RN	NA	ERROR	L	NONE	RE+RN	RE+RN	L	NA	RE	RE
IS*	RN	U	ERROR	L	NONE	D	D	L	NA	RN	RN
IS*	RE+RN	NONE	ERROR	L	NONE	NA	L	L	NA	RE+RN	RE+RN
IS*	RE+RN	L	ERROR	L	NONE	U	L	L	NA	D	D
IS*	RE+RN	RE	ERROR	L	RE	NONE	RE	L	NA	NA	L
IS*	RE+RN	RN	ERROR	L	RE	L	RE	L	NA	U	L
IS*	RE+RN	RE+RN	ERROR	L	RE	RE	RE	L	U	NONE	L
IS*	RE+RN	D	ERROR	L	RE	RN	RE+RN	L	U	L	L
IS*	RE+RN	U	ERROR	L	RE	RE+RN	RE+RN	L	U	RE	RE
IS*	D	NONE	ERROR	L	RE	D	D	L	U	RN	RN
IS*	D	L	ERROR	L	RE	NA	RE	L	U	RE+RN	RE+RN
IS*	D	RE	ERROR	L	RE	U	RE	L	U	D	D
IS*	D	RN	ERROR	L	RN	NONE	RN	L	U	NA	L
				L	RN	L	RN	L	U	U	L

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Extension SS77 or SS2000 result	LN or SS2000 result	SS77 or SS2000 result	Mets or SS2000 result	SS77 or SS2000 result	Final or SS2000 result	Extension SS77 or SS2000 result	LN or SS2000 result	SS77 or SS2000 result	Mets or SS2000 result	SS77 or SS2000 result	Final or SS2000 result	Extension SS77 or SS2000 result	LN or SS2000 result	SS77 or SS2000 result	Mets or SS2000 result	SS77 or SS2000 result	Final or SS2000 result
RE	NONE	NONE	RE	RE	RE	RE	NA	NONE	RE	RNOS	RE+RN	NONE	RNOS				
RE	NONE	L	RE	RE	RE	RE	NA	L	RE	RNOS	RE+RN	L	RNOS				
RE	NONE	RE	RE	RE	RE	RE	NA	RE	RE	RNOS	RE+RN	RE	RNOS				
RE	NONE	RN	RE+RN	RE	RE+RN	RE	NA	RN	RE+RN	RNOS	RE+RN	RN	RNOS				
RE	NONE	RE+RN	RE+RN	RE	RE+RN	RE	NA	RE+RN	RE+RN	RNOS	RE+RN	RE+RN	RNOS				
RE	NONE	D	D	RE	D	RE	NA	D	D	RNOS	RE+RN	D	D				
RE	NONE	NA	RE	RE	RE	RE	NA	NA	RE	RNOS	RE+RN	NA	RNOS				
RE	NONE	U	RE	RE	RE	RE	NA	U	RE	RNOS	RE+RN	U	RNOS				
RE	RE	NONE	RE	RE	RE	RE	U	NONE	RE	RNOS	D	NONE	D				
RE	RE	L	RE	RE	RE	RE	U	L	RE	RNOS	D	L	D				
RE	RE	RE	RE	RE	RE	RE	U	RE	RE	RNOS	D	RE	D				
RE	RE	RN	RE+RN	RE	RE+RN	RE	U	RN	RE+RN	RNOS	D	RN	D				
RE	RE	RE+RN	RE+RN	RE	RE+RN	RE	U	RE+RN	RE+RN	RNOS	D	RE+RN	D				
RE	RE	D	D	RE	D	RE	U	D	D	RNOS	D	D	D				
RE	RE	NA	RE	RE	RE	RE	U	NA	RE	RNOS	D	NA	D				
RE	RE	U	RE	RE	RE	RE	U	U	RE	RNOS	D	U	D				
RE	RN	NONE	RE+RN	RNOS	RE+RN	RNOS	NONE	NONE	RNOS	RNOS	NA	NONE	RNOS				
RE	RN	L	RE+RN	RNOS	RE+RN	RNOS	NONE	L	RNOS	RNOS	NA	L	RNOS				
RE	RN	RE	RE+RN	RNOS	RE+RN	RNOS	NONE	RE	RNOS	RNOS	NA	RE	RNOS				
RE	RN	RN	RE+RN	RNOS	RE+RN	RNOS	NONE	RN	RNOS	RNOS	NA	RN	RNOS				
RE	RN	RE+RN	RE+RN	RNOS	RE+RN	RNOS	NONE	RE+RN	RNOS	RNOS	NA	RE+RN	RNOS				
RE	RN	D	D	RNOS	D	RNOS	NONE	D	D	RNOS	NA	D	D				
RE	RN	NA	RE+RN	RNOS	RE+RN	RNOS	NONE	NA	RNOS	RNOS	NA	NA	RNOS				
RE	RN	U	RE+RN	RNOS	RE+RN	RNOS	NONE	U	RNOS	RNOS	NA	U	RNOS				
RE	RE+RN	NONE	RE+RN	RNOS	RE+RN	RNOS	RE	NONE	RNOS	RNOS	U	NONE	RNOS				
RE	RE+RN	L	RE+RN	RNOS	RE+RN	RNOS	RE	L	RNOS	RNOS	U	L	RNOS				
RE	RE+RN	RE	RE+RN	RNOS	RE+RN	RNOS	RE	RE	RNOS	RNOS	U	RE	RNOS				
RE	RE+RN	RN	RE+RN	RNOS	RE+RN	RNOS	RE	RN	RNOS	RNOS	U	RN	RNOS				
RE	RE+RN	RE+RN	RE+RN	RNOS	RE+RN	RNOS	RE	RE+RN	RNOS	RNOS	U	RE+RN	RNOS				
RE	RE+RN	D	D	RNOS	D	RNOS	RE	D	D	RNOS	U	D	D				
RE	RE+RN	NA	RE+RN	RNOS	RE+RN	RNOS	RE	NA	RNOS	RNOS	U	NA	RNOS				
RE	RE+RN	U	RE+RN	RNOS	RE+RN	RNOS	RE	U	RNOS	RNOS	U	U	RNOS				
RE	D	NONE	D	RNOS	D	RNOS	RN	NONE	RNOS	D	NONE	NONE	D				
RE	D	L	D	RNOS	D	RNOS	RN	L	RNOS	D	NONE	L	D				
RE	D	RE	D	RNOS	D	RNOS	RN	RE	RNOS	D	NONE	RE	D				
RE	D	RN	D	RNOS	D	RNOS	RN	RN	RNOS	D	NONE	RN	D				
RE	D	RE+RN	D	RNOS	D	RNOS	RN	RE+RN	RNOS	D	NONE	RE+RN	D				
RE	D	D	D	RNOS	D	RNOS	RN	D	D	D	NONE	D	D				
RE	D	NA	D	RNOS	D	RNOS	RN	NA	RNOS	D	NONE	NA	D				
RE	D	U	D	RNOS	D	RNOS	RN	U	RNOS	D	NONE	U	D				

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Extension SS77 or SS2000 result	LN or SS2000 result	SS77 or SS2000 result	Mets or SS2000 result	Final or SS2000 result	SS77 or SS2000 result	Extension SS77 or SS2000 result	LN or SS2000 result	SS77 or SS2000 result	Mets or SS2000 result	Final or SS2000 result	SS77 or SS2000 result	Extension SS77 or SS2000 result	LN or SS2000 result	SS77 or SS2000 result	Mets or SS2000 result	Final or SS2000 result	SS77 or SS2000 result	
D	RE	NONE	D	D	D	D	U	NONE	D	NA	D	NA	D	NONE	D	D	D	D
D	RE	L	D	D	D	D	U	L	D	NA	D	NA	D	L	D	D	D	D
D	RE	RE	D	D	D	D	U	RE	D	NA	D	NA	D	RE	D	D	D	D
D	RE	RN	D	D	D	D	U	RN	D	NA	D	NA	D	RN	D	D	D	D
D	RE	RE+RN	D	D	D	D	U	RE+RN	D	NA	D	NA	D	RE+RN	D	D	D	D
D	RE	D	D	D	D	D	U	D	D	NA	D	NA	D	D	D	D	D	D
D	RE	NA	D	D	D	D	U	NA	D	NA	D	NA	D	NA	D	D	D	D
D	RE	U	D	D	D	D	U	U	D	NA	D	NA	D	U	D	D	D	D
D	RN	NONE	D	D	NA	NA	NONE	NONE	U	NA	NA	NA	NA	NONE	U	D	D	U
D	RN	L	D	D	NA	NA	NONE	L	L	NA	NA	NA	NA	L	L	D	D	L
D	RN	RE	D	D	NA	NA	NONE	RE	RE	NA	NA	NA	NA	RE	RE	D	D	RE
D	RN	RN	D	D	NA	NA	NONE	RN	RN	NA	NA	NA	NA	RN	RN	D	D	RN
D	RN	RE+RN	D	D	NA	NA	NONE	RE+RN	RE+RN	NA	NA	NA	NA	RE+RN	RE+RN	D	D	RE+RN
D	RN	D	D	D	NA	NA	NONE	D	D	NA	NA	NA	NA	D	D	D	D	D
D	RN	NA	D	D	NA	NA	NONE	NA	U	NA	NA	NA	NA	NA	NA	D	D	NA
D	RN	U	D	D	NA	NA	NONE	U	U	NA	NA	NA	NA	U	U	D	D	U
D	RE+RN	NONE	D	D	NA	NA	RE	NONE	RE	NA	NA	NA	NA	U	NONE	D	D	U
D	RE+RN	L	D	D	NA	NA	RE	L	RE	NA	NA	NA	NA	U	L	D	D	L
D	RE+RN	RE	D	D	NA	NA	RE	RE	RE	NA	NA	NA	NA	U	RE	D	D	RE
D	RE+RN	RN	D	D	NA	NA	RE	RN	RE+RN	NA	NA	NA	NA	U	RN	D	D	RN
D	RE+RN	RE+RN	D	D	NA	NA	RE	RE+RN	RE+RN	NA	NA	NA	NA	U	RE+RN	D	D	RE+RN
D	RE+RN	D	D	D	NA	NA	RE	D	D	NA	NA	NA	NA	U	D	D	D	D
D	RE+RN	NA	D	D	NA	NA	RE	NA	RE	NA	NA	NA	NA	U	NA	D	D	U
D	RE+RN	U	D	D	NA	NA	RE	U	RE	NA	NA	NA	NA	U	U	D	D	U
D	D	NONE	D	D	NA	NA	RN	NONE	RN	U	NA	NA	NA	NONE	NONE	D	D	U
D	D	L	D	D	NA	NA	RN	L	RN	U	NA	NA	NA	NONE	L	D	D	L
D	D	RE	D	D	NA	NA	RN	RE	RE+RN	U	NA	NA	NA	NONE	RE	D	D	RE
D	D	RN	D	D	NA	NA	RN	RN	RN	U	NA	NA	NA	NONE	RN	D	D	RN
D	D	RE+RN	D	D	NA	NA	RN	RE+RN	RE+RN	U	NA	NA	NA	NONE	RE+RN	D	D	RE+RN
D	D	D	D	D	NA	NA	RN	D	D	U	NA	NA	NA	NONE	D	D	D	D
D	D	NA	D	D	NA	NA	RN	NA	RN	U	NA	NA	NA	NONE	NA	D	D	U
D	D	U	D	D	NA	NA	RN	U	RN	U	NA	NA	NA	NONE	U	D	D	U
D	NA	NONE	D	D	NA	NA	RE+RN	NONE	RE+RN	U	NA	NA	NA	RE	NONE	D	D	RE
D	NA	L	D	D	NA	NA	RE+RN	L	RE+RN	U	NA	NA	NA	RE	L	D	D	RE
D	NA	RE	D	D	NA	NA	RE+RN	RE	RE+RN	U	NA	NA	NA	RE	RE	D	D	RE
D	NA	RN	D	D	NA	NA	RE+RN	RN	RE+RN	U	NA	NA	NA	RE	RN	D	D	RE+RN
D	NA	RE+RN	D	D	NA	NA	RE+RN	RE+RN	RE+RN	U	NA	NA	NA	RE	RE+RN	D	D	RE+RN
D	NA	D	D	D	NA	NA	RE+RN	D	D	U	NA	NA	NA	RE	D	D	D	D
D	NA	NA	D	D	NA	NA	RE+RN	NA	RE+RN	U	NA	NA	NA	RE	NA	D	D	RE
D	NA	U	D	D	NA	NA	RE+RN	U	RE+RN	U	NA	NA	NA	RE	U	D	D	RE

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Extension SS77 or SS2000 result	LN or SS2000 result	SS77 Mets or SS2000 result	SS77 Final or SS2000 result	Extension SS77 or SS2000 result	LN or SS2000 result	SS77 Mets or SS2000 result	SS77 Final or SS2000 result
U	RN	NONE	RN	U	D	RE+RN	D
U	RN	L	RN	U	D	D	D
U	RN	RE	RE+RN	U	D	NA	D
U	RN	RN	RN	U	D	U	D
U	RN	RE+RN	RE+RN	U	NA	NONE	U
U	RN	D	D	U	NA	L	L
U	RN	NA	RN	U	NA	RE	RE
U	RN	U	RN	U	NA	RN	RN
U	RE+RN	NONE	RE+RN	U	NA	RE+RN	RE+RN
U	RE+RN	L	RE+RN	U	NA	D	D
U	RE+RN	RE	RE+RN	U	NA	NA	U
U	RE+RN	RN	RE+RN	U	NA	U	U
U	RE+RN	RE+RN	RE+RN	U	U	NONE	U
U	RE+RN	D	D	U	U	L	L
U	RE+RN	NA	RE+RN	U	U	RE	RE
U	RE+RN	U	RE+RN	U	U	RN	RN
U	D	NONE	D	U	U	RE+RN	RE+RN
U	D	L	D	U	U	D	D
U	D	RE	D	U	U	NA	U
U	D	RN	D				

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Appendix 4. Site Specific Factors

Schema	SSF1	SSF2	SSF3	SSF4	SSF5	SSF6
LipUpper	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
LipLower	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
OthLip	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
BaseTongue	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
AntTongue	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
GumUpper	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
GumLower	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
OthGum	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
FOM	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
HardPalate	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
SoftPalate	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
OthMouth	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
Buccal Mucosa	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
ParotidGland	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
Submandibular Gland	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
OthSalivary	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Schema	SSF1	SSF2	SSF3	SSF4	SSF5	SSF6
Oropharynx	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
AntEpiGlottis	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
Nasopharynx	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
Hypopharynx	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
OthPharynx	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
Esophagus	none					
Stomach	none					
SmallIntestine	none					
Colon	Carcino- embryonic Antigen (CEA)					
Rectosigmoid, Rectum	Carcino- embryonic Antigen (CEA)					
Anus	none					
Liver	Alpha Fetoprotein (AFP)	Fibrosis Score				
Gallbladder	none					
OthBiliary	none					
ExtraHepaticDucts	none					
Ampulla	none					
PancreasHead	none					
PancreasBodyTail	none					
OthPancreas	none					
OthDigestive	none					
NasalCavity	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
MiddleEar	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VII	Otn LN Group Levels
MaxillarySinus	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Schema	SSF1	SSF2	SSF3	SSF4	SSF5	SSF6
EthmoidSinus	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
OthSinus	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
GlotticLarynx	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
SupraLarynx	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
SubLarynx	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
OthLarynx	Size of LNs	Extracaps. Exten. LN for H&N	H&N LN Levels I-III	H&N LN Levels IV-V	H&N LN Levels VI-VIII	Oth LN Group Levels
Trachea	none					
Lung	none					
HeartMediastinum	none					
Pleura	Pleural Effusion					
OthRespiratory	none					
Bone	none					
Skin	none					
SkinEyelid	none					
Melanoma (of Skin, Vulva, Penis, Scrotum)	Measured Thickness (depth) Breslow's	Ulceration	Clinical Status of LN Mets	LDH		
MF	Peripheral Blood Involvement					
SoftTissue	none					
Peritoneum	none					
Breast	ERA	PRA	# Pos. Ipsilat Ax LNs	IHC of LNs	Molecular Studies Reg LNs	Size of Tumor Invasive Component
Vulva	none					
Vagina	none					
Cervix	none					
Corpus	none					
Ovary	CA-125					
FallopianTube	none					

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Schema	SSF1	SSF2	SSF3	SSF4	SSF5	SSF6
OthAdnexa	none					
OthFemaleGen	none					
Placenta	Prognostic Scoring Index					
Penis	none					
Prostate	PSA Lab Value	PSA	CS Path Extension	[PAP-obsolete] Prostate Apex Involvement*	Gleason's Prim and 2nd Patterns	Gleason's Score
Testis	AFP	HCG	LDH	Radical Orchiectomy	Size of Mets in LNs	
OthMaleGen	none					
Scrotum	none					
Kidney	none					
RenalPelvis	none					
Bladder	none					
Urethra	none					
OthUrinary	none					
Conjunctiva	none					
Melanoma Conjunctiva	Measured Thickness (depth) Breslow's					
OthEye	none					
MelanomaChoroid	Measured Thickness (depth) Breslow's					
MelanomaIrisCiliary	Measured Thickness (depth) Breslow's					
MelanomaOthEye	none					
LacrimalGland	none					
Orbit	none					
Retinoblastoma	Extension Evaluated at Enucleation					
Brain	WHO grade					
OthCNS	WHO grade					
Thyroid	Solitary vs Multifocal.I					
OthEndocrine	WHO grade					
KS	Assoc w/ HIV/AIDS					

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Schema	SSF1	SSF2	SSF3	SSF4	SSF5	SSF6
Lymphoma	Assoc w/ HIV/AIDS	Symptoms at Diagnosis	IPI score			
HemeRetic	none					
OthIIIDef	none					

* name and content of field changed as of version 1.02 (April 2005)

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Appendix 5. Histology Exclusion Groups

based on ICD-O-3 Morphology Codes

Histology Code Groupings for Collaborative Staging

Carcinomas	800-823, 8244, 8245, 8246, 8247; 825-867; 894
Carcinoids	8240, 8241, 8242, 8243, 8248, 8249
Melanomas	872-879
Sarcomas	871; 880-892; 899; 904; 912-913; 915-925; 937; 954-958
Other specified cancers	868-870; 893; 895-898; 900-903; 906-911; 926-936; 938-953
Mesotheliomas	905
Kaposi sarcoma	914
Lymphomas	959-972
Hematopoietic	973-976; 980-996; 997; 998

In the following table, 'x' in a cell means that category of cancer is excluded from AJCC staging for that site. The CS algorithm will output T-NA, N-NA, M-NA, Stage Group-NA. Conversely, an empty cell means that all histologies in that code grouping will generate (output) T, N, M, and Stage Group. A schema name marked with an asterisk (*) means that there is no TNM staging scheme in the sixth edition. For these sites, all histologies are included and only Summary Stage will be generated.

Schema	Carcinoma	Carcinoid	Melanoma	Sarcoma	Other specified cancers	Mesothelioma	Kaposi sarcoma	Lymphoma	Hematopoietic	Other exclusions
Lip: Upper; Lower; Other		x	x	x	x	x	x	x	x	
Base of Tongue		x	x	x	x	x	x	x	x	
Anterior 2/3 of Tongue		x	x	x	x	x	x	x	x	
Gum: Upper; Lower; NOS		x	x	x	x	x	x	x	x	
Floor of Mouth		x	x	x	x	x	x	x	x	
Hard Palate		x	x	x	x	x	x	x	x	
Soft Palate		x	x	x	x	x	x	x	x	
Other Mouth		x	x	x	x	x	x	x	x	
Buccal Mucosa		x	x	x	x	x	x	x	x	
Parotid Gland		x	x	x	x note 1	x	x	x	x	
Submandibular Gland		x	x	x	x note 1	x	x	x	x	
Other Salivary Gland		x	x	x	x note 1	x	x	x	x	
Tonsil, Oropharynx		x	x	x	x	x	x	x	x	
Anterior Surface of Epiglottis		x	x	x	x	x	x	x	x	
Nasopharynx		x	x	x	x	x	x	x	x	
Pyramidal Sinus; Hypopharynx		x	x	x	x	x	x	x	x	
Other Pharynx*										
Esophagus		x	x	x	x	x	x	x	x	
Stomach		x	x	x	x	x	x	x	x	
Small Intestine		x	x	x	x	x	x	x	x	
Colon		x	x	x	x	x	x	x	x	
Rectosigmoid; Rectum		x	x	x	x	x	x	x	x	
Anus		x	x	x	x	x	x	x	x	
Liver, intrahepatic ducts		x	x	x	x	x	x	x	x	
Gallbladder		x	x	x	x note 2	x	x	x	x	
Extrahepatic Ducts		x	x	x	x	x	x	x	x	
Ampulla of Vater		x	x	x	x	x	x	x	x	8013; 8041; 8246; 8247; 8574

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Schema	Carci- noma	Carcin- oid	Mela- noma	Sar- coma	Other specified cancers	Mesothe- lioma	Kaposi sarcoma	Lym- phoma	Hema- topoi- etic	Other exclu- sions
Other Biliary		x	x	x	x	x	x	x	x	
Pancreas: Head		x	x	x	x note 3	x	x	x	x	815_
Pancreas: Body, Tail		x	x	x	x note 3	x	x	x	x	815_
Other Pancreas		x	x	x	x note 3	x	x	x	x	815_
Other Digestive*										
Nasal Cavity		x	x	x	x	x	x	x	x	
Middle Ear*										
Maxillary Sinus		x	x	x	x	x	x	x	x	
Ethmoid Sinus		x	x	x	x	x	x	x	x	
Other Sinus*										
Glottic Larynx		x	x	x	x	x	x	x	x	
Supraglottic Larynx		x	x	x	x	x	x	x	x	
Subglottic Larynx		x	x	x	x	x	x	x	x	
Other Larynx		x	x	x	x	x	x	x	x	
Trachea*										
Lung		x		x	x	x	x	x	x	
Heart, Mediastinum	x	x	x		x	x	x	x	x	
Pleura	x	x	x	x	x		x	x	x	
Other Respiratory*										
Bone	x	x	x		x note 4	x	x	x	x	
Skin (Carcinoma)		x	x	x	x	x	x	x	x	
Eyelid (Carcinoma)		x	x		x	x	x	x	x	
Skin (Melanoma)	x	x		x	x	x	x	x	x	
Mycosis Fungoides	x	x	x	x	x	x	x	x note 5	x	
Soft Tissue	x	x	x		x note 6	x	x	x	x	
Retroperitoneum, Peritoneum	x	x	x		x note 6	x	x	x	x	
Breast		x	x	x	x	x	x	x	x	
Vulva		x	x	x	x	x	x	x	x	
Vagina		x	x	x	x	x	x	x	x	
Cervix		x	x	x	x	x	x	x	x	
Corpus		x	x	x	x note 7	x	x	x	x	
Ovary		x	x	x	x note 8	x	x	x	x	
Fallopian Tube		x	x	x	x	x	x	x	x	
Ligaments, Other Adnexa*										
Other Female Genital*										
Placenta	x	x	x	x	x note 9	x	x	x	x	
Penis		x	x	x	x	x	x	x	x	
Prostate		x	x	x	x	x	x	x	x	813_
Testis	x note 10	x	x	x	x note 10	x	x	x	x	
Other Male Genital*										
Scrotum			x			x	x	x	x	
Kidney		x	x	x	x	x	x	x	x	
Renal pelvis, Ureter		x	x	x	x	x	x	x	x	
Urinary Bladder		x	x	x	x	x	x	x	x	
Urethra		x	x	x	x	x	x	x	x	
Other Urinary*										
Conjunctiva (Carcinoma)		x	x	x	x	x	x	x	x	
Conjunctiva (Melanoma)	x	x		x	x	x	x	x	x	
Melanoma of uvea	x	x		x	x	x	x	x	x	
Other Eye*										
Iris, Ciliary Body (Melanoma)	x	x		x	x	x	x	x	x	
Choroid (Melanoma)	x	x		x	x	x	x	x	x	
Other Eye (Melanoma)	x	x		x	x	x	x	x	x	

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Schema	Carci-noma	Carcin-oid	Mela-noma	Sar-coma	Other specified cancers	Mesothelioma	Kaposi sarcoma	Lym-phoma	Hema-topoi-etic	Other exclu-sions
Lacrimal gland (Carcinoma)		x	x	x	x	x	x	x	x	
Orbit (Sarcoma)	x	x	x		x	x	x	x	x	
Retinoblastoma	x	x	x	x	x note 11	x	x	x	x	
Brain*										
Other CNS*										
Thyroid		x	x	x	x	x	x	x	x	
Other Endocrine*										
Kaposi Sarcoma—all sites*	x	x	x	x	x	x		x	x	
Lymphoma—all sites	x	x	x	x	x	x	x		x	
Hematopoietic, Retic*										
Other, Ill-Defined Sites*										

Note 1: For parotid gland, submandibular gland and other salivary gland, 8982 is included for TNM Staging.

Note 2: For gallbladder, 8980 is included for TNM Staging.

Note 3: For Pancreas (head, body, tail, other) 8971 is included for TNM Staging

Note 4: For bone, codes 9260-9342 are included for TNM Staging

Note 5: For mycosis fungoides and Sezary disease, all histologies other than 9700 and 9701 are excluded.

Note 6: For soft tissue and retroperitoneum/peritoneum, codes 8936 and 9473 are included for TNM Staging.

Note 7: For corpus, 8950 and 8951 are included for TNM Staging.

Note 8: For ovary, morphology codes 906-909 are included for TNM Staging.

Note 9: For placenta, 910 is included for TNM Staging.

Note 10: For testis, 859-865 and 906-910 are included for TNM Staging.

Note 11: For retinoblastoma, all histologies other than 951 are excluded.

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Index to Part I

In the text, page numbers are preceded by I-; for example, page I-49.

A	Abstracting rules	2	Clinical assessment	12
	Accessible primary sites	3, 14	Clinical information, use of	12
	Breast	3, 14	Clinical staging basis	30, 43, 49
	Lymph node involvement	3, 14	Coding rules summary	21
	Oral cavity	3, 14	Collaborative Staging Task Force	1
	Salivary gland	3, 14	Collaborative Staging web site	11
	Skin	3, 14	Colon	
	Thyroid	3, 14	CEA	51
	AFP	51	CS Lymph Nodes	35
	Aggregate tumor size	26	CS Site-Specific Factor 1	51
	AIDS/HIV	52	Site-Specific Factors	13
	AJCC Cancer Staging Manual	1, 11	Tumor size	27
	AJCC stage allowable values	69	Combination codes	12
	AJCC Stage Table	63	Composite tumor size	26
	AJCC staging not applicable	18	Computer Algorithm	4, 11, 21
	AJCC TNM staging	1, 11	Contiguous (direct) extension	
	Output from Collaborative Staging	4	CS Extension	29
	Algorithm, computer	4, 11	Corpus uteri	
	Allowable Values and Format for Collaborative Staging		Discontinuous metastasis	28
	Data Items	5	Inaccessible primary site	3
	Alpha fetoprotein	51	CS Extension	
	Ambiguous terminology	20	Allowable Values	5
	American Joint Committee on Cancer (AJCC)	1	Coding instructions	28
	Autopsy reports		In situ cancer	29
	CS Mets Eval	49	CS Extension - Pathologic Extension for prostate	55
	CS Reg Nodes Eval	43, 44	CS Lymph Nodes	
	CS Tumor Size/Ext Eval	30, 31	Allowable values	5
	use of	13, 18, 30	Coding instructions	33
			Physician staging	35
B			CS Mets At Dx	
Bladder			Allowable values	5
Inaccessible primary site	3		Coding instructions	47
Brain			None vs. unknown	33, 47
CS Site-Specific Factor 1	52		CS Mets Eval	
Regional Nodes Examined	46		Allowable values	5
Regional Nodes Positive	45		Coding instructions	49
Site-Specific Factors	13		CS Reg Nodes Eval	34
WHO grade	52		Allowable values	5
Breast			Coding instructions	43
Accessible primary site	3, 14		CS Site-Specific Factor 1	
CS Site-Specific Factor 1	51		Allowable values	5
CS Site-Specific Factor 2	53		Coding instructions	51
CS Site-Specific Factor 3	55		None vs. unknown	52
CS Site-Specific Factor 4	57		CS Site-Specific Factor 2	
CS Site-Specific Factor 5	59		Allowable values	5
CS Site-Specific Factor 6	61		Coding instructions	53
ERA	51		None vs. unknown	54
Inflammatory carcinoma	27		CS Site-Specific Factor 3	
Number of positive axillary nodes	55		Allowable values	5
PRA	53		Coding instructions	55
Regional lymph nodes	38		None vs. unknown	56
Site-Specific Factors	13		CS Site-Specific Factor 4	
Tumor size	26, 27, 61		Allowable values	5
Breslow's Measurement	51, 52		Coding instructions	57
CA-125	51		None vs. unknown	57, 58
			CS Site-Specific Factor 5	
C			Allowable values	5
Canadian Cancer Society	1		Coding instructions	59
Carbohydrate antigen 125	51		None vs. unknown	59
Carcinoembryonic Antigen	51		CS Site-Specific Factor 6	
Carcinoids histology inclusion and exclusion group	81		Allowable values	5
Carcinomas histology inclusion and exclusion group	81		Coding instructions	61
CEA	51		None vs. unknown	61, 62
Centers for Disease Control and Prevention	1		CS TS/Ext Eval	26
Central nervous system			CS Tumor Size	
CS Site-Specific Factor 1	52		Allowable values	5
Regional Nodes Examined	46		Coding instructions	25
Regional Nodes Positive	45		CS Tumor Size/Ext Eval	27, 28
WHO grade	52		Allowable values	5
Chips or pieces, tumor size	26		Coding instructions	30
CS Version 1st	4, 6, 21			
CS Version Latest	4, 6, 21		Cyst size, use of for tumor size	26
			Cystic mass size, use of for tumor size	26

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

D
Death certificate only cases 18
Derived fields, allowable values 6
Diagnostic confirmation 11
Diagram of Inputs and Outputs 7
Discontinuous metastasis 28
 Corpus uteri 28
 CS Extension 28
 Ovary 28
Disease progression 2, 12, 13
 CS Mets at DX 47
Display strings for Collaborative Staging 4
Distant metastases 29
Downstaging rule 8

E
Endocrine, other
 CS Site-Specific Factor 1 52
Enucleation for retinoblastoma 52
ERA 51
Esophagus
 Inaccessible primary site 3
 Tumor size 27
Estrogen receptor assay 51
Extension Size Table, example 64
Extra tables 63
Extracapsular extension, lymph nodes 35, 53

F
Familial polyposis 27
Fibrosis Score for liver 53
Fixed lymph nodes 34
Fruit and nuts size equivalents 66

G
General Guidelines 11
Gestational trophoblastic tumor 17
Gleason's patterns/score 59, 61
Grade, WHO 52

H
H&E 38
HCG 53
Head and neck
 CS Site-Specific Factor 1 51
 CS Site-Specific Factor 2 53
 CS Site-Specific Factor 3 55, 56
 CS Site-Specific Factor 4 57, 58
 CS Site-Specific Factor 5 59
 CS Site-Specific Factor 6 62
 Extracapsular extension, lymph nodes 53
 Lymph node levels 55, 57, 59, 61
 Regional lymph nodes 35
 Schemas included 13, 51, 53, 55, 57, 59, 61
 Site-Specific Factors 13, 36
 Size of lymph nodes 51
Head and neck lymph node levels 37, 59, 61
 Coding instructions 55, 57
Hematopoietic and reticuloendothelial system
 Histology-specific coding schema 15, 21
 Regional Nodes Examined 46
 Regional Nodes Positive 45
Hematopoietic histology inclusion and exclusion
 group 81
Hematoxylin and Eosin 38
Hierarchy of codes 12
Histologies excluded from TNM staging 11
Histology Exclusion Tables 63
Histology Inclusion and Exclusion Groups
 Names of 81
Histology-specific schemas 15, 21
HIV/AIDS 52
Human chorionic gonadotropin 53

I
IHC of lymph nodes for breast 57
Lung
 Adenopathy 34

 III-defined primary sites
 Regional Nodes Examined 46
 Regional Nodes Positive 45
Imaging studies
 CS Extension 29
 CS Lymph Nodes 34
 CS Mets Eval 50
 CS Reg Nodes Eval 44
 CS Tumor Size/Ext Eval 31
 Tumor size 26
Immunohistochemistry 39
Immunohistochemistry (IHC) of lymph nodes for
 breast 57
Immunoproliferative diseases
 Regional Nodes Examined 46
 Regional Nodes Positive 45
 Tumor size 27
Implementation date of 1/1/04 11
In situ tumor
 CS Extension 29
 CS Lymph Nodes 34
 Regional Nodes Examined 46
 Regional Nodes Positive 45
 Tumor size 26
Inaccessible primary sites 3, 14
 Bladder 3
 Corpus uteri 3
 CS Lymph Nodes 33, 34
 Esophagus 3
 Kidney 3
 Liver 3
 Lung 3
 New coding rule 14
 None vs. unknown 14
 Ovary 3
 Prostate 3
 Stomach 3
Inflammatory carcinoma 27
International Prognostic Index (IPI) Score for
 lymphoma 55
Invasive component
 Tumor size 26, 61
IPI score for lymphoma 55
Isolated tumor cells 38
ITCs 38

K
Kaposi sarcoma
 AIDS/HIV 52
 CS Site-Specific Factor 1 13, 52
 Histology-specific coding schema 15, 21
 HIV/AIDS 52
 Site-Specific Factors 13
Kaposi sarcoma histology inclusion and exclusion
 group 81
Kidney
 Inaccessible primary site 3

L
Lactate dehydrogenase 57
Langerhans cell histiocytosis, disseminated
 Tumor size 27
LDH 55, 57
Letterer-Siwe disease
 Tumor size 27
Leukemia
 Regional Nodes Examined 46
 Regional Nodes Positive 45
 Tumor size 27
Levels of head and neck lymph nodes 59
 Coding instructions 55, 57
Liver
 AFP 51
 CS Site-Specific Factor 1 51
 CS Site-Specific Factor 2 53
 Fibrosis score 53
 Inaccessible primary site 3
 Site-Specific Factors 13
 Hilar mass 34
 Inaccessible primary site 3

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

Mediastinal mass	34	Regional Nodes Examined.....	46
Tumor size	27	Regional Nodes Positive	45
Lymph nodes		Tumor size	27
Axillary, number positive for breast	55	N	
Clinical status for melanoma	55	N allowable values	68
Direct extension into.....	34	Names of Site-Specific Factors	76
Enlarged.....	34	National Cancer Institute of Canada	1
Extracapsular extension.....	35	National Cancer Institute Surveillance, Epidemiology and End Res	1
Fixed.....	34	National Cancer Registrars Association (NCRA)	1
Levels in head and neck	35, 55, 57, 61	NCRA	1
Matted.....	34	Needle biopsy, tumor size	27
Palpable.....	34	Neoadjuvant therapy	12
Shotty.....	34	CS Extension	28
Size.....	35	CS Lymph Nodes	33, 34
Size of metastasis for testis	59	CS Mets at DX	47
Visible swelling.....	34	CS Mets Eval	49, 50
Lymphadenopathy.....	34	CS Reg Nodes Eval	43, 44
Lymphoma		CS Tumor Size.....	26, 31
AIDS/HIV.....	52	CS Tumor Size/Ext Eval.....	30, 31
CS Site-Specific Factor 1	52	Regional Nodes Examined.....	46
CS Site-Specific Factor 2	53	Regional Nodes Positive	45
CS Site-Specific Factor 3	55	Tumor size	26
Histology-specific coding schema	15, 21	Nodule(s) in pericolic fat.....	35
HIV/AIDS.....	52	None vs. unknown.....	3, 14
IPI score.....	55	CS Mets at DX	33, 47
Lymph node involvement	34	CS Site-Specific Factor 1	52
Regional Nodes Examined.....	46	CS Site-Specific Factor 2	54
Regional Nodes Positive	45	CS Site-Specific Factor 3	55, 56
Site-Specific Factors	13	CS Site-Specific Factor 4	57, 58
Symptoms at diagnosis.....	53	CS Site-Specific Factor 5	59
Tumor size	27	CS Site-Specific Factor 6	61, 62
Lymphoma histology inclusion and exclusion group.....	81	NOS, Use of	8
M		CS Extension	29
M allowable values	69	CS Lymph Nodes	34
Mapping		CS Mets at DX	48
Columns in Collaborative Staging tables.....	4	CS Site-Specific Factor 3	56
How it was determined	8	CS Site-Specific Factor 4	58
Margins, positive		CS Site-Specific Factor 5	60
CS Extension	29	CS Site-Specific Factor 6	62
Mass, NOS, and lymph nodes.....	34	Use of for head and neck lymph nodes.....	36
Mast cell tumors, tumor size.....	27	Not applicable	
Matted lymph nodes	34	using code(s) for	12
Melanoma		NPCR	1
Breslow's measurement.....	51, 52	O	
Clinical status of lymph node mets.....	55	Operative assessment	12
CS Site-Specific Factor 1	51	Operative reports	
CS Site-Specific Factor 2	53	CS Reg Nodes Eval	34
CS Site-Specific Factor 3	55	CS Tumor Size/Ext Eval.....	26, 28
CS Site-Specific Factor 4	57	Oral cavity	
Histology-specific coding schema	15, 21	Accessible primary site	3, 14
LDH.....	57	Orchiectomy, radical	57
Site breakdown for	15	Other endocrine	
Site-Specific Factors	13	CS Site-Specific Factor 1	52
Thickness (depth)	51, 52	WHO grade	52
Tumor size	27	Other specified histology inclusion and exclusion group	81
Ulceration.....	53, 54	Ovary	
Melanomas histology inclusion and exclusion group.....	81	CA-125.....	51
Mesotheliomas histology inclusion and exclusion group	81	CS Site-Specific Factor 1	51
Micrometastasis	39	Discontinuous metastasis	28
Microscopic confirmation.....	11	Inaccessible primary site.....	3
Microscopic focus or foci.....	27	Site-Specific Factors	13
Tumor size	27		
Missing information, coding guideline.....	3		
Molecular studies	39		
Molecular studies of regional lymph nodes for breast.....	59		
Multiple myeloma, tumor size.....	27		
Multiple polyposis.....	27		
Multiple vs. single nodules for thyroid.....	51		
Mycosis fungoides			
CS Site-Specific Factor 1	51		
Histology-specific coding schema	15, 21		
Peripheral blood involvement.....	51		
Site-Specific Factors	13		
Myelodysplastic syndromes			
Tumor size	27		
Myeloproliferative diseases			

**Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices**

P

PAP.....57
 Pathological assessment.....12
 Pathological staging basis.....30, 43, 49
 Pathology reports
 CS Extension.....28
 CS Lymph Nodes.....33, 35
 CS Reg Nodes Eval.....34
 CS Tumor Size/Ext Eval.....26, 28
 Priority of.....12
 Size of lymph nodes.....35
 Tumor size.....25
 Peripheral blood involvement in MF.....51
 Physician staging
 CS Extension.....29
 CS Lymph Nodes.....35
 CS Mets at DX.....48
 Physician-assigned TNM.....29
 CS Extension.....29
 CS Lymph Nodes.....35
 CS Mets at DX.....48
 use of.....13
 Pieces or chips, tumor size.....26
 Placenta
 CS Site-Specific Factor 1.....51
 Prognostic Scoring Index.....51
 Regional Nodes Examined.....46
 Regional Nodes Positive.....45
 Site-Specific Factors.....13
 Pleura
 CS Site-Specific Factor 1.....51
 Pleural effusion.....51
 Site-Specific Factors.....13
 Pleural effusion.....51
 Polyp size, use of for tumor size.....26
 Polyposis, familial or multiple.....27
 PRA.....53
 Preoperative treatment.....12
 CS Extension.....28
 CS Lymph Nodes.....33, 34
 CS Mets at DX.....47
 CS Mets Eval.....49, 50
 CS Reg Nodes Eval.....43, 44
 CS Tumor Size.....26
 CS Tumor Size/Ext Eval.....30
 Regional Nodes Examined.....46
 Regional Nodes Positive.....45
 Tumor size.....26
 Presurgical treatment
 CS Tumor Size.....31
 CS Tumor Size/Ext Eval.....31
 Primary sites, inaccessible.....3
 Progesterone receptor assay.....53
 Prognostic Scoring Index for placenta.....51
 Progression of disease.....2, 12, 13
 CS Mets at DX.....47
 Prostate
 CS Extension - Pathologic Extension.....55
 CS Reg Nodes Eval.....44
 CS Site-Specific Factor 1.....51
 CS Site-Specific Factor 2.....53
 CS Site-Specific Factor 3.....55
 CS Site-Specific Factor 4.....57
 CS Site-Specific Factor 5.....59
 Gleason's patterns.....59
 Gleason's score.....61
 Inaccessible primary site.....3
 PAP.....57
 PSA.....53
 PSA lab value.....51
 Site-Specific Factors.....13
 Prostate specific antigen.....53
 Prostatic acid phosphatase.....57
 PSA.....51, 53

 Staging basis.....30, 43, 49
 CS Mets Eval.....49
 CS Reg Nodes Eval.....43, 44

R

Radical orchiectomy.....57
 Rectosigmoid
 CEA.....51
 CS Lymph Nodes.....35
 CS Site-Specific Factor 1.....51
 Tumor size.....27, 51
 Rectum
 CEA.....51
 CS Lymph Nodes.....35
 CS Site-Specific Factor 1.....51
 Site-Specific Factors.....13
 Tumor size.....27
 References.....8
 Regional Nodes Examined
 Allowable values.....5
 Coding instructions.....46
 Regional Nodes Positive
 Allowable values.....5
 Coding instructions.....45
 Residual disease
 CS Extension.....29
 Residual tumor.....26
 Reticuloendothelial system
 Histology-specific coding schema.....15, 21
 Regional Nodes Examined.....46
 Regional Nodes Positive.....45
 Retinoblastoma
 CS Site-Specific Factor 1.....52
 Extension at enucleation.....52
 Histology-specific coding schema.....15, 21
 Site-Specific Factors.....13
 Reverse Transcriptase Polymerase Chain Reaction
 RT-PCR.....39
 Rules
 Downstaging.....8
 Summary, one-page.....21
 Timing.....2

S
 Salivary gland
 Accessible primary site.....3, 14
 Sarcomas histology inclusion and exclusion group.....81
 SEER Program.....1
 SEER Summary Stage.....11
 Output from Collaborative Staging.....4
 Shotty lymph nodes.....34
 Site-Specific Extra Tables.....63
 Site-Specific Factors (SSFs)
 Brain.....13
 Breast.....13
 Colon.....13
 Head and neck.....13
 Kaposi sarcoma.....13
 Liver.....13
 Lymphoma.....13
 Melanoma.....13
 Mycosis fungoides.....13
 Names of.....76
 Ovary.....13
 Placenta.....13
 Pleura.....13
 Prostate.....13
 Purpose of.....12
 Rectum.....13
 Retinoblastoma.....13
 Schemas using.....13
 Testis.....13
 Thyroid.....13
 Size of lymph nodes.....35, 51
 Skin
 Accessible primary site.....3, 14
 Solitary vs. multifocal for thyroid.....51

 CS Tumor Size/Ext Eval.....30-32
 Start date of 1/1/04.....11

Collaborative Staging Manual and Coding Instructions Part I
Additional Tables and Appendices

Stomach		Timing rules.....	2, 12
Inaccessible primary site.....	3	TNM, physician-assigned	
Tumor size.....	27	use of.....	13
Storage codes for Collaborative Staging.....	4	Tumor markers.....	12
Summary Stage.....	11	Tumor size	
Allowable values.....	71	Aggregate.....	26
Conversion algorithm for all schemas.....	72	Breast.....	26, 27, 61
Output from Collaborative Staging.....	4	Chips or pieces.....	26
Summary Stage Table.....	63, 72	Composite.....	26
Summary Staging.....	1, 2	Descriptive.....	66
Surgery coding schemas.....	15	Fruit and nuts equivalents.....	66
Surgical observations		In situ cancer.....	26
use of.....	12	In situ component.....	26
Symptoms at diagnosis for lymphoma.....	53	Invasive component.....	26, 61
T		Melanoma.....	27
T allowable values.....	67	Microscopic focus or foci.....	27
Testis		Not applicable.....	27
AFP.....	51	Pieces or chips.....	26
CS Site-Specific Factor 1.....	51	Schemas not using for AJCC.....	17
CS Site-Specific Factor 2.....	53	Schemas requiring for AJCC.....	16
CS Site-Specific Factor 3.....	55	Unknown.....	27
CS Site-Specific Factor 4.....	57	U	
CS Site-Specific Factor 5.....	59	Ulcer size, use of for tumor size.....	26
HCG.....	53	Ulceration for melanoma.....	53
LDH.....	55	Unknown	
Radical orchiectomy.....	57	use of for head and neck lymph nodes.....	36
Site-Specific Factors.....	13	using code(s) for.....	3
Size of metastasis in lymph nodes.....	59	Unknown and ill-defined primary sites	
Text field on abstract.....	27	Regional Nodes Examined.....	46
CS Extension.....	29	Regional Nodes Positive.....	45
CS Lymph Nodes.....	35	Unknown vs. none.....	3, 14
CS Mets at DX.....	48	W	
CS Tumor Size.....	27	Web site, Collaborative Staging.....	11
Thickness (depth) of melanoma.....	51, 52	WHO grade.....	52
Thyroid		Y	
Accessible primary site.....	3, 14	Y staging basis.....	31
CS Site-Specific Factor 1.....	51		
Single vs. multiple nodules.....	51		
Site-Specific Factors.....	13		